

NEW QUANTUM BOUNDARIES FOR q -SIMPSON'S TYPE INEQUALITIES FOR CO-ORDINATED CONVEX FUNCTIONS

NECMETTIN ALP, MUHAMMAD AAMIR ALI, HÜSEYİN BUDAK, AND MEHMET ZEKİ SARİKAYA

ABSTRACT. The aim of this work is to develop quantum estimates for q -Simpson type integral inequalities for co-ordinated convex functions by using the notion of newly defined $q_1 q_2$ -derivatives and integrals. For this, we establish a new identity including quantum integrals and quantum numbers via $q_1 q_2$ -differentiable functions. After that, with the help of this equality, we achieved the results we want. The outcomes raised in this paper are extensions and generalizations of the comparable results in the literature on Simpson's inequalities for co-ordinated convex functions.

1. INTRODUCTION

Simpson's rules are well-known techniques for the numerical integration and numerical estimation of definite integrals. This method is known to be developed by Thomas Simpson (1710–1761). However, Johannes Kepler used a similar approximation about 100 years ago, so this method is also known as Kepler's rule.

Simpson's quadrature formula (Simpson's 1/3 rule) is stated as:

$$\int_{\alpha}^{\beta} \phi(x) dx \approx \frac{\beta - \alpha}{6} \left[\phi(\alpha) + 4\phi\left(\frac{\alpha + \beta}{2}\right) + \phi(\beta) \right].$$

There are a large number of estimations related to these quadrature rules in the literature, one of them is the following estimation known as Simpson's inequality:

Theorem 1. Suppose that $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is a four times continuously differentiable mapping on (α, β) , and let $\|\phi^{(4)}\|_{\infty} = \sup_{x \in (\alpha, \beta)} |\phi^{(4)}(x)| < \infty$. Then, one has the inequality

$$\left| \frac{1}{3} \left[\frac{\phi(\alpha) + \phi(\beta)}{2} + 2\phi\left(\frac{\alpha + \beta}{2}\right) \right] - \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \phi(x) dx \right| \leq \frac{1}{2880} \|\phi^{(4)}\|_{\infty} (\beta - \alpha)^4.$$

In recent years, many authors have focused on Simpson type inequalities for various classes of functions. Specifically, some mathematicians have worked on Simpson and Newton type results for convex mappings, because convexity theory is an effective and strong method for solving a great number of problems which arise within different branches of pure and applied mathematics. For example, Dragomir et al. presented new Simpson type results and their applications to quadrature formulas in numerical integration in [14]. What is more, some inequalities of Simpson type for s -convex functions are deduced by Alomari et al. in [5]. Afterwards, Sarikaya et al. observed the variants of Simpson type inequalities based on convexity in [35]. For more recent results, one can read [16, 20, 34].

On the other side, in the field of q -analysis, many studies have recently been carried out, starting with Euler owing to a vast requirement for mathematics that models quantum computing q -calculus occurred for the relationship between physics and mathematics. In different areas of mathematics, it has numerous applications such as combinatorics, number theory, basic hypergeometric functions, orthogonal polynomials, and other sciences, mechanics, the theory of relativity, and quantum theory [17–19, 21, 23]. Apparently, Euler invented this important mathematics branch. He used the q parameter in Newton's work on infinite series. Later, in a methodical manner, the q -calculus that

Key words and phrases. Simpson's inequality, q_1, q_2 -integral, quantum calculus, co-ordinated convexity, q_1, q_2 -derivatives.

2010 Mathematics Subject Classification 26D10, 26D15 26B25.

knew without limits calculus was firstly given by Jackson [17]. In 1908-1909, the general form of the q -integral and q -difference operator is defined by Jackson [21]. In 1969, for the first time Agarwal [3] defined the q -fractional derivative. In 1966-1967, Al-Salam [6] introduced a q -analog of the q -fractional integral and q -Riemann-Liouville fractional. In 2004, Rajkovic gave a definition of the Riemann-type q -integral which was generalized to Jackson q -integral. In 2013, Tariboon introduced ${}_{\alpha}D_q$ -difference operator [7]. Recently, in 2020, Bermudo et al. introduced the notion of ${}^{\beta}D_q$ derivative and integral [9].

Many well-known integral inequalities such as Hölder inequality, Hermite-Hadamard inequality, Simpson's inequality, Newton's inequality, Ostrowski inequality, Cauchy-Bunyakovsky-Schwarz inequality, Gruss inequality, Gruss-Cebysev inequality and other integral inequalities have been studied in the setup of q -calculus using the concept of classical convexity. For more results in this direction, we refer to [1, 2, 7, 8, 11, 13, 15, 17–19, 22, 25, 26, 29–33, 36, 38–43].

Inspired by this ongoing study, we establish some new quantum bounds for q -Simpson's type inequalities for q -differentiable co-ordinated convex functions. This is the primary motivation of this paper. The ideas and strategies of the paper may open new venues for further research in this field.

2. PRELIMINARIES OF q -CALCULUS AND SOME INEQUALITIES

In this section, we present some required definitions and related inequalities about q -calculus. Throughout the paper, we consider that $0 < q, q_1, q_2 < 1$, $\alpha < \beta$, $\gamma < \delta$, and $\Delta = [\alpha, \beta] \times [\gamma, \delta] \subseteq \mathbb{R}^2$.

We have to give the following notation which will be used many times in the next sections (see, [23]):

$$[n]_q = \frac{q^n - 1}{q - 1}.$$

Definition 1. [37] We consider that a function $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is continuous. Then, the q_{α} -derivative of ϕ at $\varkappa \in [\alpha, \beta]$ is defined in the following way;

$$(2.1) \quad {}_{\alpha}d_q\phi(\varkappa) = \frac{\phi(\varkappa) - \phi(q\varkappa + (1-q)\alpha)}{(1-q)(\varkappa - \alpha)}, \quad \varkappa \neq \alpha.$$

If we assume $\varkappa = \alpha$ in (2.1), we define ${}_{\alpha}d_q\phi(\alpha) = \lim_{\varkappa \rightarrow \alpha} {}_{\alpha}d_q\phi(\varkappa)$ if it exists and it is finite.

Definition 2. [37] We assume that a function $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is continuous. Then, the q_{α} -definite integral on $[\alpha, \beta]$ is defined as:

$$(2.2) \quad \int_{\alpha}^{\varkappa} \phi(\tau) {}_{\alpha}d_q\tau = (1-q)(\varkappa - \alpha) \sum_{n=0}^{\infty} q^n \phi(q^n \varkappa + (1-q^n)\alpha)$$

for $\varkappa \in [\alpha, \beta]$.

Moreover, we give the succeeding lemma which is necessary to prove the key results of this paper:

Lemma 1. We have the equality

$$\int_{\alpha}^{\beta} (\varkappa - \alpha)^a {}_{\alpha}d_q\varkappa = \frac{(\beta - \alpha)^{a+1}}{[a+1]_q}$$

for $a \in \mathbb{R} \setminus \{-1\}$.

In [7], Alp et al. established the succeeding quantum integral inequality of Hermite-Hadamard type for the convex functions in the developing of q -calculus:

Theorem 2. (q_{α} -Hermite-Hadamard inequality) We assume that a function $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is convex differentiable function on $[\alpha, \beta]$ and $0 < q < 1$. Then, we have the succeeding inequality:

$$(2.3) \quad \phi\left(\frac{q\alpha + \beta}{[2]_q}\right) \leq \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \phi(\varkappa) {}_{\alpha}d_q\varkappa \leq \frac{q\phi(\alpha) + \phi(\beta)}{[2]_q}.$$

In [7] and [30], the authors offered some estimates for the right and left hand sides of the inequality (2.3).

On the other side, Bermudo et al. gave the following new definitions and related Hermite-Hadamard type inequalities in quantum calculus:

Definition 3. [9] We consider that a function $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is continuous. Then, the q^β -derivative of ϕ at $\varkappa \in [\alpha, \beta]$ is defined in the following way;

$$(2.4) \quad {}^\beta d_q \phi(\varkappa) = \frac{\phi(\varkappa) - \phi(q\varkappa + (1-q)\beta)}{(1-q)(\varkappa - \beta)}, \quad \varkappa \neq \beta.$$

If we consider $\varkappa = \beta$ in (2.4), we define ${}_\alpha d_q \phi(\beta) = \lim_{\varkappa \rightarrow \beta} {}^\beta d_q \phi(\varkappa)$ if it exists and it is finite.

Definition 4. [9] We assume that a function $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is continuous. Then, the q^β -definite integral on $[\alpha, \beta]$ is defined as:

$$\int_{\varkappa}^{\beta} \phi(\tau) {}^\beta d_q \tau = (1-q)(\beta - \varkappa) \sum_{n=0}^{\infty} q^n \phi(q^n \varkappa + (1-q^n)\beta)$$

for $\varkappa \in [\alpha, \beta]$.

Theorem 3. (q^β -Hermite-Hadamard inequality, [9]) We assume that a function $\phi : [\alpha, \beta] \rightarrow \mathbb{R}$ is convex differentiable function on $[\alpha, \beta]$. Then, we obtain the succeeding inequality

$$(2.5) \quad \phi\left(\frac{\alpha + q\beta}{[2]_q}\right) \leq \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \phi(\varkappa) {}^\beta d_q \varkappa \leq \frac{\phi(\alpha) + q\phi(\beta)}{[2]_q}.$$

In [10], Budak offered some estimates for the right and left hand sides of the inequality (2.5).

In [28], Latif defined $q_{\alpha\gamma}$ -integral and partial q -derivatives for two variables functions as follows:

Definition 5. We suppose that a two variables function $\phi : \Delta \rightarrow \mathbb{R}$ is continuous. Then, the definite $q_{\alpha\gamma}$ -integral on Δ is defined by

$$\begin{aligned} \int_{\alpha}^{\varkappa} \int_{\gamma}^y \phi(\tau, \sigma) {}_\gamma d_{q_2} \sigma {}_\alpha d_{q_1} \tau &= (1-q_1)(1-q_2)(\varkappa - \alpha)(y - \gamma) \\ &\times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \phi(q_1^n \varkappa + (1-q_1^n)\alpha, q_2^m y + (1-q_2^m)\gamma) \end{aligned}$$

for $(\varkappa, y) \in \Delta$.

Lemma 2. [24] If the assumptions of Definition 5 hold, then

$$\begin{aligned} \int_{y_1}^y \int_{\varkappa_1}^{\varkappa} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma &= \int_{y_1}^y \int_{\alpha}^{\varkappa} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma - \int_{y_1}^y \int_{\alpha}^{\varkappa_1} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma \\ &= \int_{\gamma}^y \int_{\alpha}^{\varkappa} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma - \int_{\gamma}^{y_1} \int_{\alpha}^{\varkappa} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma \\ &\quad - \int_{\gamma}^y \int_{\alpha}^{\varkappa_1} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma + \int_{\gamma}^{y_1} \int_{\alpha}^{\varkappa_1} \phi(\tau, \sigma) {}_\alpha d_{q_1} \tau {}_\gamma d_{q_2} \sigma. \end{aligned}$$

Definition 6. [28] We consider that a two variables function $\phi : \Delta \rightarrow \mathbb{R}$ is continuous. Then, the partial q_1 -derivatives, q_2 -derivatives, and $q_1 q_2$ -derivatives at $(\varkappa, y) \in \Delta$ can be given as follows:

$$\begin{aligned} \frac{\alpha \partial_{q_1} \phi(\varkappa, y)}{\alpha \partial_{q_1} \varkappa} &= \frac{\phi(\varkappa, y) - \phi(q_1 \varkappa + (1 - q_1) \alpha, y)}{(1 - q_1)(\varkappa - \alpha)}, \quad \varkappa \neq \beta \\ \frac{\gamma \partial_{q_2} \phi(\varkappa, y)}{\gamma \partial_{q_2} y} &= \frac{\phi(\varkappa, y) - \phi(\varkappa, q_2 y + (1 - q_2) \gamma)}{(1 - q_2)(y - \gamma)}, \quad y \neq \gamma \\ \frac{\alpha, \gamma \partial_{q_1, q_2}^2 \phi(\varkappa, y)}{\alpha \partial_{q_1} \varkappa \gamma \partial_{q_2} y} &= \frac{1}{(\varkappa - \alpha)(y - \gamma)(1 - q_1)(1 - q_2)} [\phi(q_1 \varkappa + (1 - q_1) \alpha, q_2 y + (1 - q_2) \gamma) \\ &\quad + \phi(\varkappa, y) - \phi(q_1 \varkappa + (1 - q_1) \alpha, y) - \phi(\varkappa, q_2 y + (1 - q_2) \gamma)], \quad \varkappa \neq \alpha, y \neq \gamma. \end{aligned}$$

For more details related to q -integrals and derivatives for the functions of two variables (see, [28]). On the other side, Budak et al. gave the following definitions of q_α^δ , q_β^γ and $q^{\beta\delta}$ integrals:

Definition 7. [12] We suppose that a two variables function $\phi : \Delta \rightarrow \mathbb{R}$ is continuous. Then, the following q_α^δ , q_γ^β , and $q^{\beta\delta}$ integrals on Δ are defined by

$$\begin{aligned} \int_{\alpha}^{\varkappa} \int_y^{\delta} \phi(\tau, \sigma) \, {}^\delta d_{q_2} \sigma \, {}_\alpha d_{q_1} \tau &= (1 - q_2)(\delta - y) \sum_{m=0}^{\infty} q_2^m \int_{\alpha}^{\varkappa} \phi(\tau, q_2^m y + (1 - q_2^m) \delta) \, {}_\alpha d_{q_1} \tau \\ &= (1 - q_1)(1 - q_2)(\varkappa - \alpha)(\delta - y) \\ &\quad \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \phi(q_1^n \varkappa + (1 - q_1^n) \alpha, q_2^m y + (1 - q_2^m) \delta) \\ (2.6) \quad \int_{\varkappa}^{\beta} \int_{\gamma}^y \phi(\tau, \sigma) \, {}_\gamma d_{q_2} \sigma \, {}^\beta d_{q_1} \tau &= (1 - q_1)(1 - q_2)(\beta - \varkappa)(y - \gamma) \\ &\quad \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \phi(q_1^n \varkappa + (1 - q_1^n) \beta, q_2^m y + (1 - q_2^m) \gamma) \end{aligned}$$

and

$$\begin{aligned} (2.7) \quad \int_{\varkappa}^{\beta} \int_y^{\delta} \phi(\tau, \sigma) \, {}^\delta d_{q_2} \sigma \, {}^\beta d_{q_1} \tau &= (1 - q_1)(1 - q_2)(\beta - \varkappa)(\delta - y) \\ &\quad \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \phi(q_1^n \varkappa + (1 - q_1^n) \beta, q_2^m y + (1 - q_2^m) \delta) \end{aligned}$$

respectively, for $(\varkappa, y) \in \Delta$.

Theorem 4. ($q_1 q_2$ -Hölder's inequality for two variables functions, [28]) Let $\varkappa, y > 0$, $p_1 > 1$ such that $\frac{1}{p_1} + \frac{1}{r_1} = 1$. Then

$$\int_0^{\varkappa} \int_0^y |\phi(\varkappa, y) G(\varkappa, y)| d_{q_1} \varkappa d_{q_2} y \leq \left(\int_0^{\varkappa} \int_0^y |\phi(\varkappa, y)|^{p_1} d_{q_1} \varkappa d_{q_2} y \right)^{\frac{1}{p_1}} \left(\int_0^{\varkappa} \int_0^y |G(\varkappa, y)|^{r_1} d_{q_1} \varkappa d_{q_2} y \right)^{\frac{1}{r_1}}.$$

In [4], Ali et al. gave new definitions of partial $q_1 q_2$ -derivatives in the following way:

Definition 8. [4] Let a two variables function $\phi : \Delta \rightarrow \mathbb{R}$ be continuous. Then the partial q_1 -derivatives, q_2 -derivatives, and $q_1 q_2$ -derivatives at $(\varkappa, y) \in \Delta$ can be given as follows:

$$\begin{aligned}
\frac{\beta \partial_{q_1} \phi(\varkappa, y)}{\beta \partial_{q_1} \varkappa} &= \frac{\phi(q_1 \varkappa + (1 - q_1) \beta, y) - \phi(\varkappa, y)}{(1 - q_1)(\beta - \varkappa)}, \quad \varkappa \neq \beta \\
\frac{\delta \partial_{q_2} \phi(\varkappa, y)}{\delta \partial_{q_2} y} &= \frac{\phi(\varkappa, q_2 y + (1 - q_2) \delta) - \phi(\varkappa, y)}{(1 - q_2)(\delta - y)}, \quad \delta \neq y \\
\frac{\alpha \partial_{q_1}^2 \phi(\varkappa, y)}{\alpha \partial_{q_1} \varkappa \delta \partial_{q_2} y} &= \frac{1}{(\varkappa - \alpha)(\delta - y)(1 - q_1)(1 - q_2)} [\phi(q_1 \varkappa + (1 - q_1) \alpha, q_2 y + (1 - q_2) \delta) \\
&\quad - \phi(q_1 \varkappa + (1 - q_1) \alpha, y) - \phi(\varkappa, q_2 y + (1 - q_2) \delta) + \phi(\varkappa, y)], \quad \varkappa \neq \alpha, y \neq \delta, \\
\frac{\beta \partial_{q_1}^2 \phi(\varkappa, y)}{\beta \partial_{q_1} \varkappa \gamma \partial_{q_2} y} &= \frac{1}{(\beta - \varkappa)(y - \gamma)(1 - q_1)(1 - q_2)} [\phi(q_1 \varkappa + (1 - q_1) \beta, q_2 y + (1 - q_2) \gamma) \\
&\quad - \phi(q_1 \varkappa + (1 - q_1) \beta, y) - \phi(\varkappa, q_2 y + (1 - q_2) \gamma) + \phi(\varkappa, y)], \quad \varkappa \neq \beta, y \neq \gamma, \\
\frac{\beta, \delta \partial_{q_1}^2 \phi(\varkappa, y)}{\beta \partial_{q_1} \varkappa \delta \partial_{q_2} y} &= \frac{1}{(\beta - \varkappa)(\delta - y)(1 - q_1)(1 - q_2)} [\phi(q_1 \varkappa + (1 - q_1) \beta, q_2 y + (1 - q_2) \delta) \\
&\quad - \phi(q_1 \varkappa + (1 - q_1) \beta, y) - \phi(\varkappa, q_2 y + (1 - q_2) \delta) + \phi(\varkappa, y)], \quad \varkappa \neq \beta, y \neq \delta, \\
\frac{\beta \partial_{q_1} \phi(\varkappa, y)}{\beta \partial_{q_1} \varkappa} &= \frac{\phi(\varkappa, y) - \phi(q_1 \varkappa + (1 - q_1) \beta, y)}{(1 - q_1)(\varkappa - \beta)}, \quad \varkappa \neq \beta, \\
\frac{\delta \partial_{q_1} \phi(\varkappa, y)}{\delta \partial_{q_2} y} &= \frac{\phi(\varkappa, y) - \phi(\varkappa, q_2 y + (1 - q_2) \delta)}{(1 - q_2)(y - \delta)}, \quad y \neq \gamma, \\
\frac{\alpha, \gamma \partial_{q_1}^2 \phi(\varkappa, y)}{\alpha \partial_{q_1} \varkappa \gamma \partial_{q_2} y} &= \frac{1}{(\varkappa - \alpha)(y - \gamma)(1 - q_1)(1 - q_2)} [\phi(q_1 \varkappa + (1 - q_1) \alpha, q_2 y + (1 - q_2) \gamma) \\
&\quad - \phi(q_1 \varkappa + (1 - q_1) \alpha, y) - \phi(\varkappa, q_2 y + (1 - q_2) \gamma) + \phi(\varkappa, y)], \quad \varkappa \neq \alpha, y \neq \gamma.
\end{aligned}$$

3. CRUCIAL LEMMAS

In this section, we offer a new identity involving the quantum derivatives, quantum integrals, and quantum numbers. Moreover, we give some calculated quantum integrals.

Let us start with the following useful lemma.

Lemma 3. Let $\phi : \Delta \rightarrow \mathbb{R}$ be a twice partially $q_1 q_2$ -differentiable function on Δ° . If the partial $q_1 q_2$ -derivative $\frac{\beta, \delta \partial_{q_1}^2 \phi(\tau, \sigma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma}$ is continuous and integrable on Δ . Then the following identity holds for $q_1 q_2$ -integrals:

$$\begin{aligned}
(3.1) \quad & \beta, \delta \mathcal{I}_{q_1, q_2}(\phi) \\
&= (\beta - \alpha)(\delta - \gamma) \int_0^1 \int_0^1 \Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma) \frac{\beta, \delta \partial_{q_1}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma,
\end{aligned}$$

where

$$\begin{aligned}
& {}^{\beta, \delta} \mathcal{I}_{q_1, q_2}(\phi) \\
= & \frac{1}{[6]_{q_1} [6]_{q_2}} \left\{ \frac{q_1 [4]_{q_1}}{q_2} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \gamma \right) + q_1 [4]_{q_1} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \delta \right) \right. \\
& + q_1 q_2 [4]_{q_1} [4]_{q_2} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) + \frac{q_2 [4]_{q_2}}{q_1} \phi \left(\alpha, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) + q_2 [4]_{q_2} \phi \left(\beta, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) \Big\} \\
& + \frac{\phi(\alpha, \gamma) + q_1 \phi(\beta, \gamma) + q_2 \phi(\alpha, \delta) + q_1 q_2 \phi(\beta, \delta)}{q_1 q_2 [6]_{q_1} [6]_{q_2}} \\
& - \frac{1}{q_1 q_2 [6]_{q_2} (\beta - \alpha)} \int_{\alpha}^{\beta} \left[\phi(\varkappa, \gamma) + q_2^2 [4]_{q_2} \phi \left(\varkappa, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) + q_2 \phi(\varkappa, \delta) \right] {}^{\beta} d_{q_1} \varkappa \\
& - \frac{1}{q_1 q_2 [6]_{q_1} (\delta - \gamma)} \int_{\gamma}^{\delta} \left[q_1 \phi(\beta, y) + q_1^2 [4]_{q_1} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, y \right) + \phi(\alpha, y) \right] {}^{\delta} d_{q_2} y \\
& + \frac{1}{q_1 q_2 (\beta - \alpha) (\delta - \gamma)} \int_{\alpha}^{\beta} \int_{\gamma}^{\delta} \phi(\varkappa, y) {}^{\beta} d_{q_1} \varkappa {}^{\delta} d_{q_2} y
\end{aligned}$$

and

$$\Lambda_{q_1}(\tau) = \begin{cases} \tau - \frac{1}{[6]_{q_1}}, & \tau \in \left[0, \frac{1}{[2]_{q_1}}\right) \\ \tau - \frac{[5]_{q_1}}{[6]_{q_1}}, & \tau \in \left[\frac{1}{[2]_{q_1}}, 1\right], \end{cases}$$

$$\Lambda_{q_2}(\sigma) = \begin{cases} \sigma - \frac{1}{[6]_{q_2}}, & \sigma \in \left[0, \frac{1}{[2]_{q_2}}\right) \\ \sigma - \frac{[5]_{q_2}}{[6]_{q_2}}, & \sigma \in \left[\frac{1}{[2]_{q_2}}, 1\right]. \end{cases}$$

Proof. Because of the Lemma 2 and Definitions of $\Lambda_{q_1}(\tau)$ and $\Lambda_{q_2}(\sigma)$, it is easy to see that

$$\begin{aligned}
(3.2) \quad & \int_0^1 \int_0^1 \Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma) \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{{}^{\beta} \partial_{q_1} \tau {}^{\delta} \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
= & \frac{([5]_{q_2} - 1)([5]_{q_1} - 1)}{[6]_{q_1} [6]_{q_2}} \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{{}^{\beta} \partial_{q_1} \tau {}^{\delta} \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& + \frac{[5]_{q_1} - 1}{[6]_{q_1}} \int_0^{\frac{1}{[2]_{q_1}}} \int_0^1 \left(\sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right) \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{{}^{\beta} \partial_{q_1} \tau {}^{\delta} \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& + \frac{[5]_{q_2} - 1}{[6]_{q_2}} \int_0^1 \int_0^{\frac{1}{[2]_{q_2}}} \left(\tau - \frac{[5]_{q_2}}{[6]_{q_2}} \right) \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{{}^{\beta} \partial_{q_1} \tau {}^{\delta} \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& + \int_0^1 \int_0^1 \left(\tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right) \left(\sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right) \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{{}^{\beta} \partial_{q_1} \tau {}^{\delta} \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
= & I_1 + I_2 + I_3 + I_4.
\end{aligned}$$

By considering the Definition 8, we have

$$\begin{aligned}
& \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{{}^{\beta} \partial_{q_1} \tau {}^{\delta} \partial_{q_2} \sigma} \\
= & \frac{1}{(1 - q_1)(1 - q_2)(\beta - \alpha)(\delta - \gamma)\tau\sigma} [\phi(\tau q_1 \alpha + (1 - \tau q_1) \beta, \sigma q_2 \gamma + (1 - \sigma q_2) \delta) \\
& - \phi(\tau q_1 \alpha + (1 - \tau q_1) \beta, \sigma \gamma + (1 - \sigma) \delta) - \phi(\tau \alpha + (1 - \tau) \beta, \sigma q_2 \gamma + (1 - \sigma q_2) \delta) \\
& + \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)].
\end{aligned}$$

It is necessary to compute the integrals in the right side of (3.2) to conclude the proof. By using the definition of $q_1 q_2$ -integrals, we obtain that

$$\begin{aligned}
(3.3) \quad & \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
&= \frac{1}{(1-q_1)(1-q_2)(\beta-\alpha)(\delta-\gamma)} \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} \frac{1}{\tau\sigma} [\phi(\tau q_1 \alpha + (1-\tau q_1)\beta, \sigma q_2 \gamma + (1-\sigma q_2)\delta) \\
&\quad - \phi(\tau q_1 \alpha + (1-\tau q_1)\beta, \sigma\gamma + (1-\sigma)\delta) - \phi(\tau\alpha + (1-\tau)\beta, \sigma q_2 \gamma + (1-\sigma q_2)\delta) \\
&\quad + \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)] d_{q_1} \tau d_{q_2} \sigma \\
&= \frac{1}{(\beta-\alpha)(\delta-\gamma)} \left\{ \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \phi \left(\frac{q_1^{n+1}}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^{n+1}}{[2]_{q_1}} \right) \beta, \frac{q_2^{m+1}}{[2]_{q_2}} \gamma + \left(1 - \frac{q_2^{m+1}}{[2]_{q_2}} \right) \delta \right) \right. \\
&\quad - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \phi \left(\frac{q_1^{n+1}}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^{n+1}}{[2]_{q_1}} \right) \beta, \frac{q_2^m}{[2]_{q_2}} \gamma + \left(1 - \frac{q_2^m}{[2]_{q_2}} \right) \delta \right) \\
&\quad - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \phi \left(\frac{q_1^n}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^n}{[2]_{q_1}} \right) \beta, \frac{q_2^{m+1}}{[2]_{q_2}} \gamma + \left(1 - \frac{q_2^{m+1}}{[2]_{q_2}} \right) \delta \right) \\
&\quad \left. + \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \phi \left(\frac{q_1^n}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^n}{[2]_{q_1}} \right) \beta, \frac{q_2^m}{[2]_{q_2}} \gamma + \left(1 - \frac{q_2^m}{[2]_{q_2}} \right) \delta \right) \right\} \\
&= \frac{1}{(\beta-\alpha)(\delta-\gamma)} \left[\phi(\beta, \delta) - \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \delta \right) - \phi \left(\beta, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) + \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) \right].
\end{aligned}$$

For that reason, we obtain that

$$I_1 = \frac{([5]_{q_1} - 1)([5]_{q_2} - 1)}{[6]_{q_1}[6]_{q_2}(\beta-\alpha)(\delta-\gamma)} \left[\phi(\beta, \delta) - \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \delta \right) - \phi \left(\beta, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) + \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) \right].$$

Now from Definition 7, we obtain the following

$$\begin{aligned}
(3.4) \quad & \int_0^{\frac{1}{[2]_{q_1}}} \int_0^1 \sigma \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
&= \frac{1}{(1-q_1)(1-q_2)(\beta-\alpha)(\delta-\gamma)} \int_0^{\frac{1}{[2]_{q_1}}} \int_0^1 \frac{1}{\tau} [\phi(\tau q_1 \alpha + (1-\tau q_1)\beta, \sigma q_2 \gamma + (1-\sigma q_2)\delta) \\
&\quad - \phi(\tau q_1 \alpha + (1-\tau q_1)\beta, \sigma\gamma + (1-\sigma)\delta) - \phi(\tau\alpha + (1-\tau)\beta, \sigma q_2 \gamma + (1-\sigma q_2)\delta) \\
&\quad + \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)] d_{q_1} \tau d_{q_2} \sigma \\
&= \frac{1}{(\beta-\alpha)(\delta-\gamma)} \left\{ \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_2^m \left(\phi \left(\frac{q_1^{n+1}}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^{n+1}}{[2]_{q_1}} \right) \beta, q_2^{m+1} \gamma + \left(1 - q_2^{m+1} \right) \delta \right) \right) \right. \\
&\quad - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_2^m \left(\phi \left(\frac{q_1^{n+1}}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^{n+1}}{[2]_{q_1}} \right) \beta, q_2^m \gamma + \left(1 - q_2^m \right) \delta \right) \right) \\
&\quad - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_2^m \left(\phi \left(\frac{q_1^n}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^n}{[2]_{q_1}} \right) \beta, q_2^{m+1} \gamma + \left(1 - q_2^{m+1} \right) \delta \right) \right) \\
&\quad \left. + \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_2^m \left(\phi \left(\frac{q_1^n}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^n}{[2]_{q_1}} \right) \beta, q_2^m \gamma + \left(1 - q_2^m \right) \delta \right) \right) \right\}
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \sum_{m=0}^{\infty} q_2^m \left[\sum_{n=0}^{\infty} \phi \left(\frac{q_1^{n+1}}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^{n+1}}{[2]_{q_1}} \right) \beta, q_2^{m+1} \gamma + (1 - q_2^{m+1}) \delta \right) \right. \right. \\
&\quad - \sum_{n=0}^{\infty} \left(\phi \left(\frac{q_1^n}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^n}{[2]_{q_1}} \right) \beta, q_2^{m+1} \gamma + (1 - q_2^{m+1}) \delta \right) \right) \Big] \\
&\quad + \sum_{m=0}^{\infty} q_2^m \left[\sum_{n=0}^{\infty} \phi \left(\frac{q_1^n}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^n}{[2]_{q_1}} \right) \beta, q_2^m \gamma + (1 - q_2^m) \delta \right) \right. \\
&\quad \left. \left. - \sum_{n=0}^{\infty} \phi \left(\frac{q_1^{n+1}}{[2]_{q_1}} \alpha + \left(1 - \frac{q_1^{n+1}}{[2]_{q_1}} \right) \beta, q_2^m \gamma + (1 - q_2^m) \delta \right) \right] \right\} \\
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \sum_{m=0}^{\infty} q_2^m \phi (\beta, q_2^{m+1} \gamma + (1 - q_2^{m+1}) \delta) - \sum_{m=0}^{\infty} q_2^m \phi (\beta, q_2^m \gamma + (1 - q_2^m) \delta) \right. \\
&\quad + \sum_{m=0}^{\infty} q_2^m \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, q_2^m \gamma + (1 - q_2^m) \delta \right) - \sum_{m=0}^{\infty} q_2^m \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, q_2^{m+1} \gamma + (1 - q_2^{m+1}) \delta \right) \Big\} \\
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \frac{1 - q_2}{q_2} \sum_{m=0}^{\infty} q_2^m \phi (\beta, q_2^m \gamma + (1 - q_2^m) \delta) - \frac{1}{q_2} \phi (\beta, \gamma) \right. \\
&\quad \left. - \frac{1 - q_2}{q_2} \sum_{m=0}^{\infty} q_2^m \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, q_2^m \gamma + (1 - q_2^m) \delta \right) + \frac{1}{q_2} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \gamma \right) \right\} \\
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left[\frac{1}{q_2(\delta - \gamma)} \int_{\gamma}^{\delta} \phi (\beta, y) {}^{\delta}d_{q_2} y - \frac{1}{q_2(\delta - \gamma)} \int_{\gamma}^{\delta} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, y \right) {}^{\delta}d_{q_2} y \right. \\
&\quad \left. - \frac{1}{q_2} \phi (\beta, \gamma) + \frac{1}{q_2} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \gamma \right) \right].
\end{aligned}$$

By using the similar operations used in (3.3), we have

$$\begin{aligned}
(3.5) \quad & \int_0^{\frac{1}{[2]_{q_1}}} \int_0^1 \frac{\beta, {}^{\delta}d_{q_1,q_2}^2 \phi (\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left[\phi (\beta, \delta) - \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \delta \right) - \phi (\beta, \gamma) + \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \gamma \right) \right].
\end{aligned}$$

From (3.4) and (3.5), we obtain that

$$\begin{aligned}
I_2 &= \frac{[5]_{q_1} - 1}{[6]_{q_1} (\beta - \alpha)(\delta - \gamma)} \left\{ \frac{1}{q_2(\delta - \gamma)} \int_{\gamma}^{\delta} \phi (\beta, y) {}^{\delta}d_{q_2} y - \frac{1}{q_2(\delta - \gamma)} \int_{\gamma}^{\delta} \phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, y \right) {}^{\delta}d_{q_2} y \right\} \\
&\quad + \frac{[5]_{q_2} ([5]_{q_1} - 1)}{[6]_{q_1} [6]_{q_2} (\beta - \alpha)(\delta - \gamma)} \left[\phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \delta \right) - \phi (\beta, \delta) \right] \\
&\quad + \frac{[5]_{q_1} - 1}{q_2 [6]_{q_1} [6]_{q_2} (\beta - \alpha)(\delta - \gamma)} \left[\phi \left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \gamma \right) - \phi (\beta, \gamma) \right].
\end{aligned}$$

Similarly, we have

$$I_3 = \frac{[5]_{q_2} - 1}{[6]_{q_2} (\beta - \alpha)(\delta - \gamma)} \left\{ \frac{1}{q_1(\beta - \alpha)} \int_{\alpha}^{\beta} \phi (\varkappa, \delta) {}^{\beta}d_{q_1} \varkappa - \frac{1}{q_1(\beta - \alpha)} \int_{\alpha}^{\beta} \phi \left(\varkappa, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) {}^{\beta}d_{q_1} \varkappa \right\}$$

$$\begin{aligned}
& + \frac{[5]_{q_2} - 1}{q_1 [6]_{q_1} [6]_{q_2} (\beta - \alpha) (\delta - \gamma)} \left[\phi \left(\alpha, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) - \phi(\alpha, \delta) \right] \\
& + \frac{[5]_{q_1} ([5]_{q_2} - 1)}{[6]_{q_1} [6]_{q_2} (\beta - \alpha) (\delta - \gamma)} \left[\phi \left(\beta, \frac{\gamma + q_2 \delta}{[2]_{q_2}} \right) - \phi(\beta, \delta) \right].
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
(3.6) \quad & \int_0^1 \int_0^1 \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& = \frac{1}{(\beta - \alpha) (\delta - \gamma)} [\phi(\beta, \delta) - \phi(\alpha, \delta) - \phi(\beta, \gamma) + \phi(\alpha, \gamma)],
\end{aligned}$$

$$\begin{aligned}
(3.7) \quad & \int_0^1 \int_0^1 \sigma \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& = \frac{1}{(\beta - \alpha) (\delta - \gamma)} \left\{ \frac{1}{q_2 (\delta - \gamma)} \int_{\gamma}^{\delta} \phi(\beta, y) \delta d_{q_2} y - \frac{1}{q_2 (\delta - \gamma)} \int_{\gamma}^{\delta} \phi(\alpha, y) \delta d_{q_2} y \right. \\
& \quad \left. - \frac{1}{q_2} \phi(\beta, \gamma) + \frac{1}{q_2} \phi(\alpha, \gamma) \right\},
\end{aligned}$$

$$\begin{aligned}
(3.8) \quad & \int_0^1 \int_0^1 \tau \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& = \frac{1}{(\beta - \alpha) (\delta - \gamma)} \left\{ \frac{1}{q_1 (\beta - \alpha)} \int_{\alpha}^{\beta} \phi(\varkappa, \delta) \beta d_{q_1} \varkappa - \frac{1}{q_1 (\beta - \alpha)} \int_{\alpha}^{\beta} \phi(\varkappa, \gamma) \beta d_{q_1} \varkappa \right. \\
& \quad \left. - \frac{1}{q_1} \phi(\alpha, \delta) + \frac{1}{q_1} \phi(\alpha, \gamma) \right\}
\end{aligned}$$

and

$$\begin{aligned}
(3.9) \quad & \int_0^1 \int_0^1 \tau \sigma \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} d_{q_1} \tau d_{q_2} \sigma \\
& = \frac{1}{(\beta - \alpha) (\delta - \gamma)} \left\{ \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^{n+1} \alpha + (1 - q_1^{n+1}) \beta, q_2^{m+1} \gamma + (1 - q_2^{m+1}) \delta) \right. \\
& \quad - \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^{n+1} \alpha + (1 - q_1^{n+1}) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \\
& \quad - \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^{m+1} \gamma + (1 - q_2^{m+1}) \delta) \\
& \quad \left. + \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \right\}
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \frac{1}{q_1 q_2} \left[\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \right. \right. \\
&\quad - \sum_{m=0}^{\infty} q_2^m \phi(\alpha, q_2^m \gamma + (1 - q_2^m) \delta) - \sum_{n=0}^{\infty} q_1^n \phi(q_1^n \alpha + (1 - q_1^n) \beta, \gamma) + \phi(\alpha, \gamma) \Big] \\
&\quad - \frac{1}{q_1} \left[\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \right. \\
&\quad - \sum_{m=0}^{\infty} q_2^m \phi(\alpha, q_2^m \gamma + (1 - q_2^m) \delta) \Big] - \frac{1}{q_2} \left[\sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \right. \\
&\quad \left. \left. - \sum_{n=0}^{\infty} q_1^n \phi(q_1^n \alpha + (1 - q_1^n) \beta, \gamma) \right] + \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \right\} \\
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \frac{(1 - q_1)(1 - q_2)}{q_1 q_2} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} q_1^n q_2^m \phi(q_1^n \alpha + (1 - q_1^n) \beta, q_2^m \gamma + (1 - q_2^m) \delta) \right. \\
&\quad - \frac{1 - q_2}{q_1 q_2} \sum_{m=0}^{\infty} q_2^m \phi(\alpha, q_2^m \gamma + (1 - q_2^m) \delta) - \frac{1 - q_1}{q_1 q_2} \sum_{n=0}^{\infty} q_1^n \phi(q_1^n \alpha + (1 - q_1^n) \beta, \gamma) + \frac{1}{q_1 q_2} \phi(\alpha, \gamma) \Big\} \\
&= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \frac{1}{q_1 q_2 (\beta - \alpha)(\delta - \gamma)} \int_{\alpha}^{\beta} \int_{\gamma}^{\delta} \phi(\varkappa, y) {}^{\beta}d_{q_1} \varkappa {}^{\delta}d_{q_2} y \right. \\
&\quad \left. - \frac{1}{q_1 q_2 (\delta - \gamma)} \int_{\gamma}^{\delta} \phi(\alpha, y) {}^{\delta}d_{q_2} y - \frac{1}{q_1 q_2 (\beta - \alpha)} \int_{\alpha}^{\beta} \phi(\varkappa, \gamma) {}^{\beta}d_{q_1} \varkappa + \frac{1}{q_1 q_2} \phi(\alpha, \gamma) \right\}.
\end{aligned}$$

From (3.6)-(3.9), we obtain that

$$\begin{aligned}
I_4 &= \frac{1}{(\beta - \alpha)(\delta - \gamma)} \left\{ \frac{1}{q_1 q_2 (\beta - \alpha)(\delta - \gamma)} \int_{\alpha}^{\beta} \int_{\gamma}^{\delta} \phi(\varkappa, y) {}^{\beta}d_{q_1} \varkappa {}^{\delta}d_{q_2} y \right. \\
&\quad - \frac{1}{q_1 q_2 (\beta - \alpha)} \int_{\alpha}^{\beta} \phi(\varkappa, \gamma) {}^{\beta}d_{q_1} \varkappa - \frac{[5]_{q_2}}{[6]_{q_2}} \frac{1}{q_1 (\beta - \alpha)} \int_{\alpha}^{\beta} \phi(\varkappa, \delta) {}^{\beta}d_{q_1} \varkappa \\
&\quad + \frac{[5]_{q_2}}{[6]_{q_2}} \frac{1}{q_1 (\beta - \alpha)} \int_{\alpha}^{\beta} \phi(\varkappa, \gamma) {}^{\beta}d_{q_1} \varkappa - \frac{1}{q_1 q_2 (\delta - \gamma)} \int_{\gamma}^{\delta} \phi(\alpha, y) {}^{\delta}d_{q_2} y \\
&\quad - \frac{[5]_{q_1}}{[6]_{q_1}} \frac{1}{q_2 (\delta - \gamma)} \int_{\gamma}^{\delta} \phi(\beta, y) {}^{\delta}d_{q_2} y + \frac{[5]_{q_1}}{[6]_{q_1}} \frac{1}{q_2 (\delta - \gamma)} \int_{\gamma}^{\delta} \phi(\alpha, y) {}^{\delta}d_{q_2} y \\
&\quad \left. + \frac{\phi(\alpha, \gamma)}{q_1 q_2 [6]_{q_1} [6]_{q_2}} + \frac{[5]_{q_1} [5]_{q_2}}{[6]_{q_1} [6]_{q_2}} \phi(\beta, \delta) + \frac{[5]_{q_2}}{q_1 [6]_{q_1} [6]_{q_2}} \phi(\alpha, \delta) + \frac{[5]_{q_1}}{q_2 [6]_{q_1} [6]_{q_2}} \phi(\beta, \gamma) \right\}.
\end{aligned}$$

Now, the calculated integrals $(I_1) - (I_4)$ yield

$$\begin{aligned}
&I_1 + I_2 + I_3 + I_4 \\
&= \frac{1}{[6]_{q_1} [6]_{q_2} (\beta - \alpha)(\delta - \gamma)} \left\{ \frac{q_1 [4]_{q_1}}{q_2} \phi\left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \gamma\right) + q_1 [4]_{q_1} \phi\left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \delta\right) \right. \\
&\quad + q_1 q_2 [4]_{q_1} [4]_{q_2} \phi\left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, \frac{\gamma + q_2 \delta}{[2]_{q_2}}\right) + \frac{q_2 [4]_{q_2}}{q_1} \phi\left(\alpha, \frac{\gamma + q_2 \delta}{[2]_{q_2}}\right) + q_2 [4]_{q_2} \phi\left(\beta, \frac{\gamma + q_2 \delta}{[2]_{q_2}}\right) \Big\} \\
&\quad + \frac{\phi(\alpha, \gamma) + q_1 \phi(\beta, \gamma) + q_2 \phi(\alpha, \delta) + q_1 q_2 \phi(\beta, \delta)}{q_1 q_2 [6]_{q_1} [6]_{q_2} (\beta - \alpha)(\delta - \gamma)} \\
&\quad - \frac{1}{q_1 q_2 [6]_{q_2} (\beta - \alpha)^2 (\delta - \gamma)} \int_{\alpha}^{\beta} \left[\phi(\varkappa, \gamma) + q_2^2 [4]_{q_2} \phi\left(\varkappa, \frac{\gamma + q_2 \delta}{[2]_{q_2}}\right) + q_2 \phi(\varkappa, \delta) \right] {}^{\beta}d_{q_1} \varkappa
\end{aligned}$$

$$\begin{aligned} & -\frac{1}{q_1 q_2 [6]_{q_1} (\beta - \alpha) (\delta - \gamma)^2} \int_\gamma^\delta \left[q_1 \phi(\beta, y) + q_1^2 [4]_{q_1} \phi\left(\frac{\alpha + q_1 \beta}{[2]_{q_1}}, y\right) + \phi(\alpha, y) \right] {}^\delta d_{q_2} y \\ & + \frac{1}{q_1 q_2 (\beta - \alpha)^2 (\delta - \gamma)^2} \int_\alpha^\beta \int_\gamma^\delta \phi(\varkappa, y) {}^\beta d_{q_1} \varkappa {}^\delta d_{q_2} y \end{aligned}$$

and multiplying the resultant one with $(\beta - \alpha)(\delta - \gamma)$, we obtain the desirable equality (3.1) which accomplishes the proof. \square

Remark 1. Under the given conditions of Lemma 3 with $q_1, q_2 \rightarrow 1^-$, we obtain the succeeding identity

$$\begin{aligned} (3.10) \quad & \frac{\phi\left(\frac{\alpha+\beta}{2}, \gamma\right) + \phi\left(\frac{\alpha+\beta}{2}, \delta\right) + 4\phi\left(\frac{\alpha+\beta}{2}, \frac{\gamma+\delta}{2}\right) + \phi\left(\alpha, \frac{\gamma+\delta}{2}\right) + \phi\left(\beta, \frac{\gamma+\delta}{2}\right)}{9} \\ & + \frac{\phi(\alpha, \gamma) + \phi(\alpha, \delta) + \phi(\beta, \gamma) + \phi(\beta, \delta)}{36} \\ & - \frac{1}{6(\beta - \alpha)} \int_\alpha^\beta \left[\phi(\varkappa, \gamma) + 4\phi\left(\varkappa, \frac{\gamma+\delta}{2}\right) + \phi(\varkappa, \delta) \right] d\varkappa \\ & - \frac{1}{6(\delta - \gamma)} \int_\gamma^\delta \left[\phi(\alpha, y) + 4\phi\left(\frac{\alpha+\beta}{2}, y\right) + \phi(\beta, y) \right] dy \\ & + \frac{1}{(\beta - \alpha)(\delta - \gamma)} \int_\alpha^\beta \int_\gamma^\delta \phi(\varkappa, y) d\varkappa dy \\ = & (\beta - \alpha)(\delta - \gamma) \int_0^1 \int_0^1 \Lambda(\tau) \Lambda(\sigma) \frac{\partial^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\partial \tau \partial \sigma} d\tau d\sigma \end{aligned}$$

where

$$\Lambda(\tau) = \begin{cases} \tau - \frac{1}{6}, & \tau \in [0, \frac{1}{2}] \\ \tau - \frac{5}{6}, & \tau \in [\frac{1}{2}, 1] \end{cases}$$

and

$$\Lambda(\sigma) = \begin{cases} \sigma - \frac{1}{6}, & \sigma \in [0, \frac{1}{2}] \\ \sigma - \frac{5}{6}, & \sigma \in [\frac{1}{2}, 1] \end{cases}$$

which is proved by Özdemir et al. in [34, Lemma 1].

For reasons of brevity, let us prove another lemma that we will use frequently in our transactions:

Lemma 4. The following q -integrals hold:

$$(3.11) \quad A_1(q) = \int_0^{\frac{1}{[2]_q}} \tau \left| \tau - \frac{1}{[6]_q} \right| d_q \tau = q^2 \left(\frac{2 [2]_q^2 + q [3]_q [6]_q^2}{[2]_q^3 [3]_q [6]_q^3} \right),$$

$$(3.12) \quad A_2(q) = \int_0^{\frac{1}{[2]_q}} (1 - \tau) \left| \tau - \frac{1}{[6]_q} \right| d_q \tau = q \left(\frac{2 [2]_q^2 [3]_q [6]_q - 2q [2]_q^2 + (-1 + q^3 + q^4) [3]_q [6]_q^2}{[2]_q^3 [3]_q [6]_q^3} \right),$$

$$\begin{aligned} (3.13) \quad A_3(q) &= \int_{\frac{1}{[2]_q}}^1 \tau \left| \tau - \frac{[5]_q}{[6]_q} \right| d_q \tau \\ &= \frac{2q^2 [2]_q^2 [5]_q^3 - (q + 3q^2 + 5q^3 + 6q^4 + 4q^5 + 2q^6) [6]_q^2}{[2]_q^3 [3]_q [6]_q^3}, \end{aligned}$$

$$(3.14) \quad A_4(q) = \int_{\frac{1}{[2]_q}}^1 \left| \tau - \frac{[5]_q}{[6]_q} \right| (1-\tau) d_q \tau \\ = \frac{2 \left[(q^5 + q^6 - 1) [3]_q + [2]_q [5]_q \right] [5]_q^2}{[2]_q [3]_q [6]_q^3} + \frac{1 + [2]_q^2 - q [2]_q [5]_q}{[2]_q^3 [6]_q} - \frac{[2]_q^3 + 1}{[2]_q^3 [3]_q},$$

$$(3.15) \quad A_5(q) = \int_0^{\frac{1}{[2]_q}} \left| \tau - \frac{1}{[6]_q} \right| d_q \tau = \frac{2q [2]_q^2 + [6]_q^2 - [2]_q^2 [6]_q}{[2]_q^3 [6]_q^2},$$

$$(3.16) \quad A_6(q) = \int_{\frac{1}{[2]_q}}^1 \left| \tau - \frac{[5]_q}{[6]_q} \right| d_q \tau = \frac{\left(1 + [2]_q^2 \right) [6]_q^2 - \left(1 + [7]_q \right) [2]_q^2 [5]_q}{[2]_q^3 [6]_q^2}.$$

Proof. Since $\frac{1}{[2]_q} \geq \frac{1}{[6]_q}$ for $q \in (0, 1)$, we have the following equality:

$$\begin{aligned} A_1(q) &= \int_0^{\frac{1}{[2]_q}} \tau \left| \tau - \frac{1}{[6]_q} \right| d_q \tau \\ &= \int_0^{\frac{1}{[6]_q}} \tau \left(\frac{1}{[6]_q} - \tau \right) d_q \tau + \int_{\frac{1}{[6]_q}}^{\frac{1}{[2]_q}} \tau \left(\tau - \frac{1}{[6]_q} \right) d_q \tau \\ &= \int_0^{\frac{1}{[6]_q}} \left(\frac{\tau}{[6]_q} - \tau^2 \right) d_q \tau + \int_{\frac{1}{[6]_q}}^{\frac{1}{[2]_q}} \tau \left(\tau - \frac{1}{[6]_q} \right) d_q \tau \\ &= \left(\frac{\tau^2}{[2]_q [6]_q} - \frac{\tau^3}{[3]_q} \right)_0^{\frac{1}{[6]_q}} + \left(\frac{\tau^3}{[3]_q} - \frac{\tau^2}{[2]_q [6]_q} \right)_{\frac{1}{[6]_q}}^{\frac{1}{[2]_q}} \\ &= q^2 \frac{2 [2]_q^2 + q [3]_q [6]_q^2}{[2]_q^3 [3]_q [6]_q^3}. \end{aligned}$$

Similarly, the q -integrals (3.12)-(3.16) can be obtained and the proof is completed. \square

4. SOME NEW $q_1 q_2$ -SIMPSON'S TYPE INEQUALITIES

In this section, we prove some new quantum boundaries for quantum Simpson's inequalities using the Lemma 3.

Theorem 5. *We assume that the conditions of Lemma 3 are satisfy. Then, we obtain the succeeding inequality provided that $\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau, \sigma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|$ is convex on Δ*

$$(4.1) \quad \left| \beta, \delta \mathcal{I}_{q_1, q_2}(\phi) \right| \leq (\beta - \alpha) (\delta - \gamma) \left[(A_1(q_1) + A_3(q_1)) (A_1(q_2) + A_3(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right. \\ + (A_1(q_1) + A_3(q_1)) (A_2(q_2) + A_4(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \\ + (A_2(q_1) + A_4(q_1)) (A_1(q_2) + A_3(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \\ \left. + (A_2(q_1) + A_4(q_1)) (A_2(q_2) + A_4(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right].$$

Proof. By taking the modulus of the identity in Lemma 3, because of the properties of the modulus, we find that

$$(4.2) \quad |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \leq (\beta - \alpha)(\delta - \gamma) \int_0^1 \int_0^1 |\Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma)| \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| d_{q_1} \tau d_{q_2} \sigma.$$

Using the convexity of $\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|$, the inequality (4.2) becomes

$$(4.3) \quad |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \leq (\beta - \alpha)(\delta - \gamma) \int_0^1 \Lambda_{q_2}(\sigma) \left[\int_0^1 \Lambda_{q_1}(\tau) \left\{ \tau \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| + (1-\tau) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right\} d_{q_1} \tau \right] d_{q_2} \sigma.$$

Now, we compute the integrals appeared in the right side of the inequality (4.3)

$$\begin{aligned} & \int_0^1 \Lambda_{q_1}(\tau) \left\{ \tau \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| + (1-\tau) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right\} d_{q_1} \tau \\ = & \int_0^{\frac{1}{[2]_q}} \tau \left| \tau - \frac{1}{[6]_q} \right| \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| d_{q_1} \tau \\ & + \int_0^{\frac{1}{[2]_q}} (1-\tau) \left| \tau - \frac{1}{[6]_q} \right| \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| d_{q_1} \tau \\ & + \int_{\frac{1}{[2]_q}}^1 \tau \left| \tau - \frac{[5]_q}{[6]_q} \right| \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| d_{q_1} \tau \\ & + \int_0^{\frac{1}{[2]_q}} (1-\tau) \left| \tau - \frac{[5]_q}{[6]_q} \right| \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| d_{q_1} \tau. \end{aligned}$$

From (3.11)-(3.14), we obtain that

$$\begin{aligned} & \int_0^1 \Lambda_{q_1}(\tau) \left\{ \tau \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| + (1-\tau) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right\} d_{q_1} \tau \\ = & \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| (A_1(q_1) + A_3(q_1)) \\ & + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| (A_2(q_1) + A_4(q_1)). \end{aligned}$$

Thus, we have

$$\begin{aligned}
& |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \\
\leq & (\beta - \alpha)(\delta - \gamma) \int_0^1 \Lambda_{q_2}(\sigma) \left[\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| (A_1(q_1) + A_3(q_1)) \right. \\
& \quad \left. + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| (A_2(q_1) + A_4(q_1)) \right] d_{q_2} \sigma \\
\leq & (\beta - \alpha)(\delta - \gamma) \int_0^1 \Lambda_{q_2}(\sigma) \left[\left\{ \sigma \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| + (1-\sigma) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right. \right. \\
& \quad \times (A_1(q_1) + A_3(q_1)) \} + \left\{ \sigma \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| + (1-\sigma) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right. \\
& \quad \times (A_2(q_1) + A_4(q_1)) \} \left. \right] d_{q_2} \sigma \\
= & (\beta - \alpha)(\delta - \gamma) (A_1(q_1) + A_3(q_1)) \left[\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_0^{\frac{1}{[2]_q}} \sigma \left| \sigma - \frac{1}{[6]_q} \right| d_{q_2} \sigma \right. \\
& \quad + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_0^{\frac{1}{[2]_q}} (1-\sigma) \left| \sigma - \frac{1}{[6]_q} \right| d_{q_2} \sigma + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_{\frac{1}{[2]_q}}^1 \sigma \left| \sigma - \frac{[5]_q}{[6]_q} \right| d_{q_2} \sigma \\
& \quad + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_{\frac{1}{[2]_q}}^1 (1-\sigma) \left| \sigma - \frac{[5]_q}{[6]_q} \right| d_{q_2} \sigma \left. \right] \\
& \quad + (\beta - \alpha)(\delta - \gamma) (A_2(q_1) + A_4(q_1)) \left[\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_0^{\frac{1}{[2]_q}} \sigma \left| \sigma - \frac{1}{[6]_q} \right| d_{q_2} \sigma \right. \\
& \quad + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_0^{\frac{1}{[2]_q}} (1-\sigma) \left| \sigma - \frac{1}{[6]_q} \right| d_{q_2} \sigma + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_{\frac{1}{[2]_q}}^1 \sigma \left| \sigma - \frac{[5]_q}{[6]_q} \right| d_{q_2} \sigma \\
& \quad + \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \int_{\frac{1}{[2]_q}}^1 (1-\sigma) \left| \sigma - \frac{[5]_q}{[6]_q} \right| d_{q_2} \sigma \left. \right].
\end{aligned}$$

From (3.11)-(3.14), we have

$$\begin{aligned}
|\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \leq & (\beta - \alpha)(\delta - \gamma) \left[(A_1(q_1) + A_3(q_1))(A_1(q_2) + A_3(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right. \\
& + (A_1(q_1) + A_3(q_1))(A_2(q_2) + A_4(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \\
& + (A_2(q_1) + A_4(q_1))(A_1(q_2) + A_3(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \\
& \left. + (A_2(q_1) + A_4(q_1))(A_2(q_2) + A_4(q_2)) \left| \frac{\beta, \delta \partial_{q_1, q_2} \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right| \right].
\end{aligned}$$

Hence, the proof is completed. \square

Remark 2. Under the given conditions of Theorem 5 with $q_1, q_2 \rightarrow 1^-$, we obtain the succeeding inequality

$$\begin{aligned}
(4.4) \quad & \left| \frac{\phi\left(\frac{\alpha+\beta}{2}, \gamma\right) + \phi\left(\frac{\alpha+\beta}{2}, \delta\right) + 4\phi\left(\frac{\alpha+\beta}{2}, \frac{\gamma+\delta}{2}\right) + \phi\left(\alpha, \frac{\gamma+\delta}{2}\right) + \phi\left(\beta, \frac{\gamma+\delta}{2}\right)}{9} \right. \\
& + \frac{\phi(\alpha, \gamma) + \phi(\alpha, \delta) + \phi(\beta, \gamma) + \phi(\beta, \delta)}{36} \\
& - \frac{1}{6(\beta-\alpha)} \int_{\alpha}^{\beta} \left[\phi(\varkappa, \gamma) + 4\phi\left(\varkappa, \frac{\gamma+\delta}{2}\right) + \phi(\varkappa, \delta) \right] d\varkappa \\
& - \frac{1}{6(\delta-\gamma)} \int_{\gamma}^{\delta} \left[\phi(\alpha, y) + 4\phi\left(\frac{\alpha+\beta}{2}, y\right) + \phi(\beta, y) \right] dy \\
& \left. + \frac{1}{(\beta-\alpha)(\delta-\gamma)} \int_{\alpha}^{\beta} \int_{\gamma}^{\delta} \phi(\varkappa, y) d\varkappa dy \right| \\
& \leq \frac{25(\beta-\alpha)(\delta-\gamma)}{72} \left[\left| \frac{\partial^2 \phi(\alpha, \gamma)}{\partial \tau \partial \sigma} \right| + \left| \frac{\partial^2 \phi(\alpha, \delta)}{\partial \tau \partial \sigma} \right| + \left| \frac{\partial^2 \phi(\beta, \gamma)}{\partial \tau \partial \sigma} \right| + \left| \frac{\partial^2 \phi(\beta, \delta)}{\partial \tau \partial \sigma} \right| \right]
\end{aligned}$$

which is shown by Özdemir et al. in [34, Theorem 3].

Theorem 6. We assume that the conditions of Lemma 3 hold. If $\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau, \sigma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p$ is convex on Δ for some $p > 1$ and $\frac{1}{r} + \frac{1}{p} = 1$. Then, we obtain the succeeding inequality

$$\begin{aligned}
(4.5) \quad & |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \\
& \leq (\beta-\alpha)(\delta-\gamma) \frac{\left\{ q_1^r [5]_{q_1}^r + q_1^{5r} ([2]_{q_1}^{r+1} - 1) \right\}^{\frac{1}{r}} \left\{ q_2^r [5]_{q_2}^r + q_2^{5r} ([2]_{q_2}^{r+1} - 1) \right\}^{\frac{1}{r}}}{[2]_{q_1}^2 [2]_{q_2}^2 [6]_{q_1} [6]_{q_2} [r+1]_{q_1}^{\frac{1}{r}} [r+1]_{q_2}^{\frac{1}{r}}} \\
& \times \left(\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + q_2 \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right. \\
& \left. + q_1 \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + q_1 q_2 \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right)^{\frac{1}{p}}.
\end{aligned}$$

Proof. Applying the well-known Hölder's inequality for $q_1 q_2$ -integrals to the integrals in the right side of (4.2), it is found that

$$\begin{aligned}
(4.6) \quad & |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \\
& \leq (\beta-\alpha)(\delta-\gamma) \left[\left(\int_0^1 \int_0^1 |\Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma)|^r d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{r}} \right. \\
& \times \left. \left(\int_0^1 \int_0^1 \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1-\tau)\beta, \sigma \gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{p}} \right].
\end{aligned}$$

By applying the convexity of $\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau, \sigma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^{p_1}$, the inequality (4.6) becomes

$$(4.7) \quad \begin{aligned} & |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \\ & \leq (\beta - \alpha)(\delta - \gamma) \left[\left(\int_0^1 \int_0^1 |\Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma)|^r d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{r}} \right. \\ & \quad \times \left(\int_0^1 \int_0^1 \left[\tau \sigma \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + \tau(1-\sigma) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right. \right. \\ & \quad \left. \left. + (1-\tau)\sigma \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + (1-\tau)(1-\sigma) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right] d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{p}}. \end{aligned}$$

Now, for $0 \leq \tau, \sigma \leq 1$, we know $\tau - \frac{1}{[6]_{q_1}} \leq \tau - \frac{\tau}{[6]_{q_1}}$, $\tau - \frac{[5]_{q_1}}{[6]_{q_1}} \leq \tau - \frac{\tau [5]_{q_1}}{[6]_{q_1}}$, and this yields

$$\begin{aligned} & \int_0^1 \int_0^1 |\Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma)|^r d_{q_1} \tau d_{q_2} \sigma \\ & = \int_0^1 \int_0^{\frac{1}{[2]_{q_2}}} |\Lambda_{q_1}(\tau)|^r \left| \sigma - \frac{1}{[6]_{q_2}} \right|^r d_{q_2} \sigma d_{q_1} \tau \\ & \quad + \int_0^1 \int_{\frac{1}{[2]_{q_2}}}^1 |\Lambda_{q_1}(\tau)|^r \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right|^r d_{q_2} \sigma d_{q_1} \tau \\ & = \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{1}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{1}{[6]_{q_2}} \right|^r d_{q_2} \sigma \\ & \quad + \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{1}{[6]_{q_2}} \right|^r d_{q_2} \sigma \\ & \quad + \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{1}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_{\frac{1}{[2]_{q_2}}}^1 \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right|^r d_{q_2} \sigma d_{q_1} \tau \\ & \quad + \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_{\frac{1}{[2]_{q_2}}}^1 \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right|^r d_{q_2} \sigma d_{q_1} \tau \\ & \leq \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{\tau}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{\sigma}{[6]_{q_2}} \right|^r d_{q_2} \sigma \\ & \quad + \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{\tau [5]_{q_1}}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{\sigma}{[6]_{q_2}} \right|^r d_{q_2} \sigma \\ & \quad + \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{\tau}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_{\frac{1}{[2]_{q_2}}}^1 \left| \sigma - \frac{\sigma [5]_{q_2}}{[6]_{q_2}} \right|^r d_{q_2} \sigma \\ & \quad + \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{\tau [5]_{q_1}}{[6]_{q_1}} \right|^r d_{q_1} \tau \int_{\frac{1}{[2]_{q_2}}}^1 \left| \sigma - \frac{\sigma [5]_{q_2}}{[6]_{q_2}} \right|^r d_{q_2} \sigma \\ & = \left| 1 - \frac{1}{[6]_{q_1}} \right|^r \left| 1 - \frac{1}{[6]_{q_2}} \right|^r \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} \tau^r \sigma^r d_{q_2} \sigma d_{q_1} \tau \\ & \quad + \left| 1 - \frac{[5]_{q_1}}{[6]_{q_1}} \right|^r \left| 1 - \frac{1}{[6]_{q_2}} \right|^r \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} \tau^r \sigma^r d_{q_2} \sigma d_{q_1} \tau \end{aligned}$$

$$\begin{aligned}
& + \left| 1 - \frac{1}{[6]_{q_1}} \right|^r \left| 1 - \frac{[5]_{q_2}}{[6]_{q_2}} \right|^r \int_0^{\frac{1}{[2]_{q_1}}} \tau^r \sigma^r d_{q_2} \sigma d_{q_1} \tau \\
& + \left| 1 - \frac{[5]_{q_1}}{[6]_{q_1}} \right|^r \left| 1 - \frac{[5]_{q_2}}{[6]_{q_2}} \right|^r \int_{\frac{1}{[2]_{q_1}}}^1 \tau^r \sigma^r d_{q_2} \sigma d_{q_1} \tau \\
= & \frac{1}{[2]_{q_1}^{r+1} [2]_{q_2}^{r+1} [6]_{q_1}^r [6]_{q_2}^r [r+1]_{q_1} [r+1]_{q_2}} \\
& \times \left\{ q_1^r q_2^r [5]_{q_1}^r [5]_{q_2}^r + q_1^{5r} q_2^r [5]_{q_2}^r ([2]_{q_1}^{r+1} - 1) + q_1^r q_2^{5r} [5]_{q_1}^r ([2]_{q_2}^{r+1} - 1) \right. \\
& \left. + q_1^{5r} q_2^{5r} ([2]_{q_1}^{r+1} - 1) ([2]_{q_2}^{r+1} - 1) \right\} \\
= & \frac{\left\{ q_1^r [5]_{q_1}^r + q_1^{5r} ([2]_{q_1}^{r+1} - 1) \right\} \left\{ q_2^r [5]_{q_2}^r + q_2^{5r} ([2]_{q_2}^{r+1} - 1) \right\}}{[2]_{q_1}^{r+1} [2]_{q_2}^{r+1} [6]_{q_1}^r [6]_{q_2}^r [r+1]_{q_1} [r+1]_{q_2}}
\end{aligned}$$

and

$$\begin{aligned}
(4.8) \quad & \left(\int_0^1 \int_0^1 |\Lambda_{q_1}(\tau) \Lambda_{q_2}(\sigma)|^r d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{r}} \\
= & \frac{\left\{ q_1^r [5]_{q_1}^r + q_1^{5r} ([2]_{q_1}^{r+1} - 1) \right\}^{\frac{1}{r}} \left\{ q_2^r [5]_{q_2}^r + q_2^{5r} ([2]_{q_2}^{r+1} - 1) \right\}^{\frac{1}{r}}}{[2]_{q_1}^{\frac{r+1}{r}} [2]_{q_2}^{\frac{r+1}{r}} [6]_{q_1} [6]_{q_2} [r+1]_{q_1}^{\frac{1}{r}} [r+1]_{q_2}^{\frac{1}{r}}}.
\end{aligned}$$

Additionally, if we apply the concept of Lemma 1 for $\alpha = 0$ to the above quantum integrals, we attain

$$(4.9) \quad \int_0^1 \int_0^1 \tau \sigma d_{q_1} \tau d_{q_2} \sigma = \left(\int_0^1 \tau d_{q_1} \tau \right) \left(\int_0^1 \sigma d_{q_2} \sigma \right) = \frac{1}{[2]_{q_1} [2]_{q_2}},$$

$$(4.10) \quad \int_0^1 \int_0^1 \tau (1 - \sigma) d_{q_1} \tau d_{q_2} \sigma = \frac{q_2}{[2]_{q_1} [2]_{q_2}},$$

$$(4.11) \quad \int_0^1 \int_0^1 (1 - \tau) \sigma d_{q_1} \tau d_{q_2} \sigma = \frac{q_1}{[2]_{q_1} [2]_{q_2}},$$

$$(4.12) \quad \int_0^1 \int_0^1 (1 - \tau) (1 - \sigma) d_{q_1} \tau d_{q_2} \sigma = \frac{q_1 q_2}{[2]_{q_1} [2]_{q_2}}.$$

By substituting the calculated integrals (4.8) and (4.9)-(4.12) in (4.7), we obtain the desired inequality (4.5) which finishes the proof. \square

Theorem 7. We suppose that the assumptions of Lemma 3 hold. If $\left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau, \sigma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p$ is convex on Δ for some $p \geq 1$. Then, we obtain the succeeding inequality

$$\begin{aligned}
(4.13) \quad & |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \\
& \leq (\beta - \alpha)(\delta - \gamma) \left[A_5^{1-\frac{1}{p}}(q_1) A_5^{1-\frac{1}{p}}(q_2) \right. \\
& \quad \times \left\{ A_1(q_1) \left(A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \right. \\
& \quad + A_2(q_1) \left(A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \left. \right\}^{\frac{1}{p}} \\
& \quad + A_5^{1-\frac{1}{p}}(q_1) A_6^{1-\frac{1}{p}}(q_2) \\
& \quad \times \left\{ A_1(q_1) \left(A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \right. \\
& \quad + A_2(q_1) \left(A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \left. \right\}^{\frac{1}{p}} \\
& \quad + A_6^{1-\frac{1}{p}}(q_1) A_5^{1-\frac{1}{p}}(q_2) \\
& \quad \times \left\{ A_3(q_1) \left(A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \right. \\
& \quad + A_4(q_1) \left(A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \left. \right\}^{\frac{1}{p}} \\
& \quad + A_6^{1-\frac{1}{p}}(q_1) A_6^{1-\frac{1}{p}}(q_2) \\
& \quad \times \left\{ A_3(q_1) \left(A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \right. \\
& \quad + A_4(q_1) \left(A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) \left. \right\}^{\frac{1}{p}} \left. \right].
\end{aligned}$$

Proof. Applying the well-known power mean inequality for $q_1 q_2$ -integrals to the integrals in the right side of (4.2), it is found that

$$\begin{aligned}
(4.14) \quad & |\beta, \delta \mathcal{I}_{q_1, q_2}(\phi)| \\
& \leq (\beta - \alpha)(\delta - \gamma) \left[\left(\int_0^{\frac{1}{[2]_{q_2}}} \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{1}{[6]_{q_1}} \right| \left| \sigma - \frac{1}{[6]_{q_2}} \right| d_{q_1} \tau d_{q_2} \sigma \right)^{1-\frac{1}{p}} \right. \\
& \quad \times \left(\int_0^{\frac{1}{[2]_{q_2}}} \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{1}{[6]_{q_1}} \right| \left| \sigma - \frac{1}{[6]_{q_2}} \right| \right. \\
& \quad \times \left. \left. \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1 - \tau) \beta, \sigma \gamma + (1 - \sigma) \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{p}} \right]
\end{aligned}$$

$$\begin{aligned}
& + \left(\int_0^{\frac{1}{[2]_{q2}}} \int_{\frac{1}{[2]_{q1}}}^1 \left| \tau - \frac{1}{[6]_{q1}} \right| \left| \sigma - \frac{[5]_{q2}}{[6]_{q2}} \right| d_{q1} \tau d_{q2} \sigma \right)^{1-\frac{1}{p}} \\
& \times \left(\int_0^{\frac{1}{[2]_{q2}}} \int_{\frac{1}{[2]_{q1}}}^1 \left| \tau - \frac{1}{[6]_{q1}} \right| \left| \sigma - \frac{[5]_{q2}}{[6]_{q2}} \right| \right. \\
& \quad \times \left. \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p d_{q1} \tau d_{q2} \sigma \right)^{\frac{1}{p}} \\
& + \left(\int_{\frac{1}{[2]_{q2}}}^1 \int_0^{\frac{1}{[2]_{q1}}} \left| \tau - \frac{[5]_{q1}}{[6]_{q1}} \right| \left| \sigma - \frac{1}{[6]_{q2}} \right| d_{q1} \tau d_{q2} \sigma \right)^{1-\frac{1}{p}} \\
& \times \left(\int_{\frac{1}{[2]_{q2}}}^1 \int_0^{\frac{1}{[2]_{q1}}} \left| \tau - \frac{[5]_{q1}}{[6]_{q1}} \right| \left| \sigma - \frac{1}{[6]_{q2}} \right| \right. \\
& \quad \times \left. \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p d_{q1} \tau d_{q2} \sigma \right)^{\frac{1}{p}} \\
& + \left(\int_{\frac{1}{[2]_{q2}}}^1 \int_{\frac{1}{[2]_{q1}}}^1 \left| \tau - \frac{[5]_{q1}}{[6]_{q1}} \right| \left| \sigma - \frac{[5]_{q2}}{[6]_{q2}} \right| d_{q1} \tau d_{q2} \sigma \right) \\
& \times \left(\int_{\frac{1}{[2]_{q2}}}^1 \int_{\frac{1}{[2]_{q1}}}^1 \left| \tau - \frac{[5]_{q1}}{[6]_{q1}} \right| \left| \sigma - \frac{[5]_{q2}}{[6]_{q2}} \right| \right. \\
& \quad \times \left. \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p d_{q1} \tau d_{q2} \sigma \right)^{\frac{1}{p}} \Big].
\end{aligned}$$

By applying the convexity of $\left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\tau, \sigma)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p$, we have

$$\begin{aligned}
(4.15) \quad & \left(\int_0^{\frac{1}{[2]_{q2}}} \int_0^{\frac{1}{[2]_{q1}}} \left| \tau - \frac{1}{[6]_{q1}} \right| \left| \sigma - \frac{1}{[6]_{q2}} \right| d_{q1} \tau d_{q2} \sigma \right)^{1-\frac{1}{p}} \\
& \times \left(\int_0^{\frac{1}{[2]_{q2}}} \int_0^{\frac{1}{[2]_{q1}}} \left| \tau - \frac{1}{[6]_{q1}} \right| \left| \sigma - \frac{1}{[6]_{q2}} \right| \right. \\
& \quad \times \left. \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p d_{q1} \tau d_{q2} \sigma \right)^{\frac{1}{p}} \\
& \leq \left(\int_0^{\frac{1}{[2]_{q1}}} \left| \tau - \frac{1}{[6]_{q1}} \right| d_{q1} \tau \int_0^{\frac{1}{[2]_{q2}}} \left| \sigma - \frac{1}{[6]_{q2}} \right| d_{q2} \sigma \right)^{1-\frac{1}{p}} \\
& \quad \left[\int_0^{\frac{1}{[2]_{q2}}} \left| \sigma - \frac{1}{[6]_{q2}} \right| \left\{ \int_0^{\frac{1}{[2]_{q1}}} \left| \tau - \frac{1}{[6]_{q1}} \right| \left(\tau \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p \right. \right. \right. \\
& \quad \left. \left. \left. + (1-\tau) \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p \right) d_{q1} \tau \right\} d_{q2} \sigma \right]^{\frac{1}{p}} \\
& = A_5^{1-\frac{1}{p}}(q_1) A_5^{1-\frac{1}{p}}(q_2) \left[A_1(q_1) \int_0^{\frac{1}{[2]_{q2}}} \left| \sigma - \frac{1}{[6]_{q2}} \right| \left| \frac{\beta, \delta \partial_{q1,q2}^2 \phi(\alpha, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q1} \tau \delta \partial_{q2} \sigma} \right|^p d_{q2} \sigma \right]^p
\end{aligned}$$

$$\begin{aligned}
& + A_2(q_1) \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{1}{[6]_{q_2}} \right| \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p d_{q_2} \sigma \Bigg]^{\frac{1}{p}} \\
\leq & A_5^{1-\frac{1}{p}}(q_1) A_5^{1-\frac{1}{p}}(q_2) \left[A_1(q_1) \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{1}{[6]_{q_2}} \right| \left(\sigma \left| \frac{\beta, d \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau d \partial_{q_2} \sigma} \right|^p \right. \right. \\
& \left. \left. + (1-\sigma) \left| \frac{\beta, d \partial_{q_1, q_2}^2 \phi(\alpha, d)}{\beta \partial_{q_1} \tau d \partial_{q_2} \sigma} \right|^p \right) d_{q_2} \sigma + A_2(q_1) \int_0^{\frac{1}{[2]_{q_2}}} \left| \sigma - \frac{1}{[6]_{q_2}} \right| \right. \\
& \times \left(\sigma \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + (1-\sigma) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right) d_{q_2} \sigma \Bigg]^{\frac{1}{p}} \\
= & A_5^{1-\frac{1}{p}}(q_1) A_5^{1-\frac{1}{p}}(q_2) \left[A_1(q_1) \left\{ A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right. \\
& \left. + A_2(q_1) \left\{ A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right]^{\frac{1}{p}}.
\end{aligned}$$

By applying the similar operations, we obtain that

$$\begin{aligned}
(4.16) \quad & \left(\int_0^{\frac{1}{[2]_{q_2}}} \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{1}{[6]_{q_1}} \right| \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right| d_{q_1} \tau d_{q_2} \sigma \right)^{1-\frac{1}{p}} \\
& \times \left(\int_0^{\frac{1}{[2]_{q_2}}} \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{1}{[6]_{q_1}} \right| \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right| \right. \\
& \times \left. \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{p}} \\
\leq & A_5^{1-\frac{1}{p}} A_6^{1-\frac{1}{p}} \left[A_1(q_1) \left\{ A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right. \\
& \left. + A_2(q_1) \left\{ A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right]^{\frac{1}{p}},
\end{aligned}$$

$$\begin{aligned}
(4.17) \quad & \left(\int_{\frac{1}{[2]_{q_2}}}^1 \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right| \left| \sigma - \frac{1}{[6]_{q_2}} \right| d_{q_1} \tau d_{q_2} \sigma \right)^{1-\frac{1}{p}} \\
& \times \left(\int_{\frac{1}{[2]_{q_2}}}^1 \int_0^{\frac{1}{[2]_{q_1}}} \left| \tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right| \left| \sigma - \frac{1}{[6]_{q_2}} \right| \right. \\
& \times \left. \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau\alpha + (1-\tau)\beta, \sigma\gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p d_{q_1} \tau d_{q_2} \sigma \right)^{\frac{1}{p}} \\
\leq & A_6^{1-\frac{1}{p}} A_5^{1-\frac{1}{p}} \left[A_3(q_1) \left\{ A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right. \\
& \left. + A_4(q_1) \left\{ A_1(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_2(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right]^{\frac{1}{p}}
\end{aligned}$$

and

$$\begin{aligned}
 (4.18) \quad & \left(\int_{\frac{1}{[2]_{q_2}}}^1 \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right| \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right| d_{q_1} \tau d_{q_2} \sigma \right) \\
 & \times \left[\int_{\frac{1}{[2]_{q_2}}}^1 \int_{\frac{1}{[2]_{q_1}}}^1 \left| \tau - \frac{[5]_{q_1}}{[6]_{q_1}} \right| \left| \sigma - \frac{[5]_{q_2}}{[6]_{q_2}} \right| \right. \\
 & \quad \times \left. \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\tau \alpha + (1-\tau)\beta, \sigma \gamma + (1-\sigma)\delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p d_{q_1} \tau d_{q_2} \sigma \right]^{\frac{1}{p}} \\
 & \leq A_6^{1-\frac{1}{p}} A_6^{1-\frac{1}{p}} \left[A_3(q_1) \left\{ A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\alpha, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right. \\
 & \quad \left. + A_4(q_1) \left\{ A_3(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \gamma)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p + A_4(q_2) \left| \frac{\beta, \delta \partial_{q_1, q_2}^2 \phi(\beta, \delta)}{\beta \partial_{q_1} \tau \delta \partial_{q_2} \sigma} \right|^p \right\} \right]^{\frac{1}{p}}.
 \end{aligned}$$

From (4.14)-(4.18), we obtain the desired inequality and the proof is ended. \square

5. CONCLUSION

In this research, we have proved a new integral identity involving quantum integrals and quantum numbers. We have proved some new Simpson's inequalities for $q_1 q_2$ -differentiable co-ordinated convex functions using the newly derived equality. It is also shown that the results presented in this research transformed into some classical results by taking the limits $q_1, q_2 \rightarrow 1^-$ in the main results. It is an interesting and new problem that the upcoming researchers may use the techniques of this research and prove Newton's inequalities and similar inequalities for different kinds of convexities in their future work.

6. ACKNOWLEDGEMENTS

The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

AVAILABILITY OF DATA AND MATERIALS

Not applicable.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHOR'S CONTRIBUTIONS

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

REFERENCES

- [1] M. A. Ali, H. Budak, Z. Zhang, and H. Yildirim, *Some new Simpson's type inequalities for co-ordinated convex functions in quantum calculus*, Mathematical Methods in the Applied Sciences, <https://doi.org/10.1002/mma.7048>.
- [2] M. A. Ali, H. Budak, M. Abbas, and Y.-M. Chu, *Quantum Hermite-Hadamard type inequalities for functions whose second q^b -derivatives absolute value are convex*, Advances in Difference Equation, In press, 2020.
- [3] R. Agarwal, "A propos d'une note de m. pierre humbert," Comptes rendus de l'Academie des Sciences, vol. 236, no. 21, pp. 2031–2032, 1953.
- [4] M. A. Ali, H. Kalsoom, H. Budak, M. Z. Sarikaya, and Y.-M. Chu, *On some new trapezoidal type inequalities for the functions of two variables via quantum calculus*, available at: <https://www.researchgate.net/publication/344469986>.
- [5] M. Alomari, M. Darus, and S. S. Dragomir, *New inequalities of Simpson's type for σ -convex functions with applications*, RGMIA Res. Rep. Coll., vol. 12, no. 4, 2009.

- [6] W. Al-Salam, *Some fractional q -integrals and q -derivatives*, Proceedings of the Edinburgh Mathematical Society, vol. 15, no. 2, pp. 135–140, 1966/1967.
- [7] N. Alp, M. Z. Sarikaya, M. Kunt and I. Iscan, *q -Hermite Hadamard inequalities and quantum estimates for midpoint type inequalities via convex and quasi-convex functions*, Journal of King Saud University–Science (2018) 30, 193–203.
- [8] N. Alp and M. Z. Sarikaya, *Quantum Hermite-Hadamard's type inequalities for co-ordinated convex functions*, Applied Mathematics E-Notes, 20(2020), 341-356.
- [9] S. Bermudo, P. Kórus and J. N. Valdés, *On q -Hermite–Hadamard inequalities for general convex functions*, Acta Mathematica Hungarica, 1-11, 2020.
- [10] H. Budak, *Some trapezoid and midpoint type inequalities for newly defined quantum integrals*, Proyecciones Journal of Mathematics, in press.
- [11] H. Budak, S. Erden, and M. A. Ali, *Simpson and Newton type inequalities for convex functions via newly defined quantum integrals*, Mathematical Methods in the Applied Sciences (2020).
- [12] H. Budak, M. A. Ali, and M. Tarhanaci, *Some New Quantum Hermite–Hadamard-Like Inequalities for Coordinated Convex Functions*, Journal of Optimization Theory and Applications (2020): 1-12.
- [13] H. Budak, M. A. Ali, and T. Tunç, *Quantum Ostrowski type integral inequalities for functions of two variables*, Mathematical Methods in the Applied Sciences, In press, 2020.
- [14] S. S. Dragomir, R. P. Agarwal, P. Cerone, *On Simpson's inequality and applications*, J. Inequal. Appl. 5 (2000) 533–579.
- [15] S. Erden, S. Iftikhar, M. R. Delavar, P. Kumam, P. Thounthong, and W. Kumam, *On generalizations of some inequalities for convex functions via quantum integrals*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas 114 (2020): 1-15.
- [16] S. Erden, S. Iftikhar, P. Kumam, and M. U. Awan, *Some Newton's like inequalities with applications*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas 114, no. 4 (2020): 1-13.
- [17] T. Ernst, *The History of Q -Calculus And New Method*. Sweden: Department of Mathematics, Uppsala University, 2000.
- [18] T. Ernst, *A Comprehensive Treatment of q -Calculus*, Springer Basel (2012).
- [19] H. Gauchman, *Integral inequalities in q -calculus*, Comput. Math. Appl. 47 (2004) 281-300.
- [20] S. Iftikhar, S. Erden, P. Kumam, and M. U. Awan, *Local fractional Newton's inequalities involving generalized harmonic convex functions*, Advances in Difference Equations 2020 (2020): 1-14.
- [21] F. H. Jackson, *On a q -definite integrals*, Quarterly J. Pure Appl. Math. 41 (1910) 193-203.
- [22] S. Jhanthanam, T. Jessada , N. Sotiris K., and N. Kamsing, *On q -Hermite-Hadamard Inequalities for Differentiable Convex Functions*, Mathematics 7, no. 7 (2019): 632.
- [23] V. Kac and P. Cheung *Quantum calculus*, Springer (2001).
- [24] M. Kunt, M. A. Latif, İ. İşcan, S. S. Dragomir, *Quantum Hermite-Hadamard type inequality and some estimates of quantum midpoint type inequalities for double integrals*, Sigma J. Eng. Nat. Sci, 37(1), 2019, 207-223.
- [25] M. A. Khan, M. Noor, E. R. Nwaeze, and Y-M. Chu , *Quantum Hermite–Hadamard inequality by means of a Green function*, Advances in Difference Equations 2020, no. 1 (2020): 1-20.
- [26] M. Kunt, İ. İmdat, N. Alp, and M. Z. Sarikaya, *(p, q)-Hermite–Hadamard inequalities and (p, q) -estimates for midpoint type inequalities via convex and quasi-convex functions*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas 112, no. 4 (2018): 969-992.
- [27] H. Kalsoom, J. D. Wu, S. Hussain and M. A. Latif, *Simpson's type inequalities for co-ordinated convex functions on quantum calculus*. Symmetry, 11(6), 2019, p.768.
- [28] M. A. Latif, S. S. Dragomir, E. Momoniat, *Some q -analogues of Hermite-Hadamard inequality of functions of two variables on finite rectangles in the plane*, Journal of King Saud University-Science(2017)29,263-273.
- [29] W. Liu and Z. Hefeng, *Some quantum estimates of Hermite-Hadamard inequalities for convex functions*, Journal of Applied Analysis and Computation, 7(2), 501-522.
- [30] M. A. Noor, K. I. Noor, M. U. Awan, *Some quantum estimates for Hermite-Hadamard inequalities*, Appl. Math. Comput. 251 (2015) 675–679.
- [31] M. A. Noor, K. I. Noor, M. U. Awan, *Some quantum integral inequalities via preinvex functions*, Appl. Math. Comput. 269 (2015) 242–251.
- [32] M. Noor, K. Noor, and M. Awan, *Quantum Ostrowski inequalities for q -differentiable convex functions*, J. Math. Inequal, 2016, 10(4), 1013-1018.
- [33] E. R. Nwaeze, and A. M. Tameru, *New parameterized quantum integral inequalities via η -quasiconvexity*, Advances in Difference Equations 2019, no. 1 (2019): 425.
- [34] M. E. Özdemir, A. O. Akdemir, H. Kavurmacı and M. Avci, *On the Simpson's inequality for co-ordinated convex functions*, Turkish Journal of Analysis and Number Theory, 2014, Vol. 2, No. 5, 165-169.
- [35] M. Z. Sarikaya, E. Set and M. E. Özdemir, *On new inequalities of Simpson's type for convex functions*, RGMIA Res. Rep. Coll. 13(2) (2010), Article2.
- [36] W. Sudsutad, S. K. Ntouyas, J. Tariboon, *Quantum integral inequalities for convex functions*, J. Math. Inequal. 9 (3) (2015) 781–793.
- [37] J. Tariboon, S. K. Ntouyas, *Quantum calculus on finite intervals and applications to impulsive difference equations*, Adv. Difference Equ. 282 (2013) 1-19.
- [38] M. Vivas-Cortez, R. Liko, A. Kashuri, J. E. Hernández Hernández, *New Quantum Estimates of Trapezium-Type Inequalities for Generalized ϕ -Convex Functions*, Mathematics, 2019, 7(11), 1-19.

- [39] M. Vivas-Cortez, R. Liko, A. Kashuri, J. E. Hernández Hernández, *Quantum Estimates of Ostrowski Inequalities for Generalized ϕ -Convex Functions* Symmetry, 2019, 11(12), 1-16.
- [40] M. Vivas-Cortez, R. Liko, A. Kashuri, J. E. Hernández Hernández, *Some New q -Integral Inequalities Using Generalized Quantum Montgomery Identity via Preinvex Functions*, Symmetry, 2020, 12(4), 1-15.
- [41] M. Vivas-Cortez, A. Kashuri, R. Liko, J. E. Hernández Hernández, *Quantum Trapezium-Type Inequalities Using Generalized f -Convex Functions* Axioms, 2020, 9(1), 1-14.
- [42] M. Vivas-Cortez, M. A. Ali, A. Kashuri, I.B. Sial, and Z. Zhang, *Some New Newton's Type Integral Inequalities for Co-Ordinated Convex Functions in Quantum Calculus*, Symmetry 12, no. 9 (2020): 1476.
- [43] H. Zhuang, W. Liu, J. Park, *Some quantum estimates of Hermite-Hadamard inequalities for quasi-convex functions*, Miskolc Math. Notes, 2016, 17(2).

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY
E-mail address: placenn@gmail.com

JIANGSU KEY LABORATORY FOR NSLSCS, SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING, 210023, CHINA.

E-mail address: mahr.muhammad.aamir@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY
E-mail address: hsyn.budak@gmail.com

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY
E-mail address: sarikayamz@gmail.com