

Gastroschisis

JULIEN SAADA, MD,*†§|| JEAN-FRANÇOIS OURY, MD,*‡§
EDITH VUILLARD, MD,* JEAN GUIBOURDENCHE, PhD,††
PASCAL DE LAGUSIE, MD, PhD,‡§||**
GHISLAINE STERKERS, MD, PhD,‡‡ JOSEPH P. BRUNER, MD,¶
and DOMINIQUE LUTON, MD, PhD*‡§||

*Département de Périnatologie, Maternité de l'Hôpital Robert Debré (AP-HP), Paris, France; †Fédération de Gynécologie-obstétrique, Service d'Echographie et de Diagnostic Anténatal, Hôpital Paule de Viguier, Toulouse, France; ‡Université Paris VII (UFR Lariboisière, Saint Louis), Paris, France; §Unité de recherche EA3102, France; ||Ecole de Chirurgie du Fer à Moulin (AP-HP), Paris, France; ¶Department of Obstetrics and Gynecology, Vanderbilt University Medical Center, Nashville, Tennessee; **Service de chirurgie Pédiatrique, Hôpital la Timone (AP-HM), Marseille, France; ††Biochimie-Hormonologie, Hôpital Robert Debré (AP-HP), Paris, France; and ‡‡Laboratoire d'immunologie, Hôpital Robert Debré (AP-HP), Paris, France

Introduction

Gastroschisis is defined as an abdominal wall defect that involves herniation of the gut and occasionally other abdominal organs such as the spleen and liver. It occurs mostly to the right side of the umbilicus and is characterized by the absence of a membrane covering the prolapsed organs with bowel loops floating in the amniotic fluid.

It is now commonly diagnosed during the prenatal period by means of sonography and the overall prognosis is excellent with a 90% survival rate.^{1,2} Morbidity depends on associated prematurity, intrauterine growth retardation, bowel atresia, and perivisceritis

resulting from contact with the amniotic fluid. The pathogenesis of gastroschisis and secondary bowel lesions are not fully understood. The management of gastroschisis is also still debated because there is no large prospective study concerning the mode, time, and place of delivery, or the type and time of initial surgery during the neonatal period.

Recent advances in the understanding of the elements responsible for the bowel damages such as digestive compounds and associated inflammation in the amniotic fluid may induce changes in therapeutic strategy, including prenatal treatment using amnioexchange.

Correspondence: Dominique Luton, MD, PhD, Département de Périnatologie, Maternité de l'Hôpital Robert Debré (AP-HP), 48 bvd Sérurier, 75019, Paris, France.
E-mail: dominique.luton@rdb.ap-hop-paris.fr

Prevalence and Epidemiology

The total prevalence of gastroschisis, including live births, stillbirths, and terminations

of pregnancy, is approximately one to 4 in 10000 live births¹⁻⁶ with a sex ratio approximately 1 (Table 1). Although the total prevalence varies among geographic areas,^{2,7,8,9} studies of several international registries or referral centers report a global increase of gastroschisis during recent years.^{1,2,7,10} No etiology has yet been implicated in this global increase, and prospective studies are warranted to identify potential etiologies.

Several maternal risk factors have been identified in association with gastroschisis. Goldbaum et al¹¹ found an increased risk from maternal age less than 25 years old with the highest risk in mothers less than 20 years old (odds ratio, 4.1). Werler et al reported that relative risks in 20- to 24-year-old and less than 20-year-old mother groups were, respectively, 5, 4, and 16.¹² More recently, several studies have reported similar results, confirming that young maternal age is a major risk factor for fetal gastroschisis.^{1,2,7,13} It occurs more frequently in primiparas^{2,13,14} and is associated with low socioeconomic status, poor maternal education, and drug abuse (tobacco, alcohol, and cocaine).^{12,15,16} Other teratogens have been implicated and are reported in Table 2. It has been suggested that these substances may have vasoreactive effects during embryonic development.

Some isolated associations have been described with aneuploidy, trisomy 13, 18, 21, and monosomy 22,¹⁷ and some familial cases have also been reported.¹⁸⁻²⁶ The role of genetic, epigenetic, and environmental factors is not yet clearly established.²⁷

TABLE 1. Increasing Incidence of Gastroschisis

	Period of Study	Incidence per 10000 Live Births
Baerg et al	1985-1990/1996-2000	1.85/4.06
Penman et al	1987/1995	1.60/4.40
Rankin et al	1986/1996	1.48/4.72
Kazaura et al	1968/1998	0.50/2.90
Houglund et al	1971/2002	0.36/3.92

TABLE 2. Potential Teratogens Implicated in Gastroschisis and Epidemiologic Associations

Potential teratogens
Tobacco
Alcohol
Cocaine
Aspirin
Ibuprofen
Pseudoephedrine
Acetaminophen
Phenylpropanolamine
Epidemiologic associations
Young maternal age and short interval between menarche and first pregnancy
Low socioeconomic status
Poor maternal education

Pathogenesis of Gastroschisis and Bowel Lesions

The pathogenesis of gastroschisis is not clearly understood; the most cited mechanism is the premature regression, by the fifth or sixth weeks of gestation, of the right omphalomesenteric artery or of the right umbilical vein.^{28,29} This premature regression leads to a failure of mesodermal components of the abdominal wall; intestinal malrotation is almost constantly associated with gastroschisis. Others consider that gastroschisis may be related to an early in utero rupture of an umbilical cord hernia.³⁰

Several studies in animal models yielded a better understanding of associated bowel lesions. Their cause seems to be both chemical and mechanical.³¹⁻³⁵ Bowel contact with the amniotic fluid is associated with a sterile inflammation of the bowel wall resulting in a fibrous coat called perivisceritis. Other bowel lesions of increasing gravity such as ischemia, perforation, stenosis, atresia, partial or complete necrosis may be present and affect morbidity, mortality, and prognosis.³⁶⁻⁴⁰

Burc et al⁴¹ found that, in cases of gastroschisis, intraamniotic values of total protein and ferritin levels were increased and displayed a chronic inflammatory profile. In another study,⁴² IL8 also displayed increased levels together with an acute inflammatory exudate composed predominantly of

neutrophil polymorphs and mononuclear cells (Table 3). Because of the occurrence of meconium stained amniotic fluid, some authors have suggested that gastrointestinal waste products might be implicated in this in utero inflammatory reaction.^{41,43,44}

Most bowel histologic studies have been performed on resection samples of atretic bowel from initial or iterative surgery or performed quickly after birth on necroscopic samples; they revealed the presence of edema, fibrin deposition, fibrosis, capillary proliferation, cellular infiltration of epithelial cells, and macrophages in the bowel wall.^{45,46} These results have also been described in animal models.⁴⁷⁻⁵⁰ In a rat model, Yu et al⁵¹ have demonstrated that late injection of dexamethasone into the amniotic fluid of fetuses with gastroschisis had beneficial effects on bowel wall thickening, normalized total DNA, induced cellular proliferation, and decreased apoptosis. In 2000, Correia-Pinto⁵² demonstrated in another rat model of gastroschisis that perivisceritis was dependent on the presence of meconium. It has also been demonstrated that exchanging amnioallantoic fluid decreased bowel lesion intensity,⁵³ which was coincident with a decrease in gastrointestinal waste products, whereas urinary waste products did not vary.

Recently, it has been suggested that in utero defecation is a physiological event during normal fetal life and increases in the last weeks of pregnancy.⁵⁴ This has been supported by the presence of intestinal enzymes in the amniotic fluid of normal pregnancies.^{55,56} Ciftci et al⁵⁴ have suggested that meconium-stained amniotic fluid is not only related to meconium passage, but perhaps to

impaired clearance of amniotic fluid; digestive compounds present in the amniotic fluid could thus be responsible for the chemical mechanism leading to perivisceritis.^{48,57} The exact impact of amniotic fluid on the peritoneal surface of the bowel wall is not yet completely known; however, exposure of the fetal pleural surface to amniotic fluid induces similar changes seen in the bowel wall in cases of gastroschisis.⁵⁸

The inextensible parietal defect may induce a mechanical compression on the superior mesenteric artery or vein and be responsible for ischemic complications or venous congestion. Occlusive and subocclusive complications may also occur and be responsible for in utero vomiting, abnormal defecation,⁵⁹ and decreased clearance of amniotic fluid together with bowel hypoperistaltism.⁵⁴

Some studies have demonstrated that bowel lesions are associated with downregulation of various genes involved in amino acid and glucose absorption⁶⁰⁻⁶³ in experimental models of gastroschisis and with fetal protein loss.⁶⁴ Other studies have demonstrated that overexpression of nitrous oxide synthase⁶⁵⁻⁶⁸ is associated with bowel hypoperistaltism.^{60,69}

Prenatal Diagnosis and Sonography

Prenatal diagnosis of gastroschisis relies on sonography. Gastroschisis, in contrast to omphalocele, is not associated with an increased risk of chromosomal abnormalities or associated extragastrointestinal malformations. Some studies have produced contradictory results; although Rankin et al have reported a malformation rate of 5.3% associated with gastroschisis and Roland et al a rate of 5.5%, the EUROCAT study has reported a rate of 20.8% of associated malformations. The most common extragastrointestinal malformations associated with gastroschisis are reported in Table 4.^{1,2}

Third-trimester sonography is not accurately able to identify partial necrosis and secondary atresia, but in a retrospective

TABLE 3. Chronic Inflammation in Amniotic Fluid

Inflammatory exudate	+
IL6	↑
IL8	↑
IL1β	—
TNFα	—
Ferritin	↑

TABLE 4. Malformations Associated With Gastroschisis

Hydronephrosis
Arthrogryposis
Hypoplastic gallbladder
Meckel diverticulum
Oligo-anhydramnios (intrauterine growth restriction)

study of 45 patients with a prenatal diagnosis of gastroschisis, Japaraj et al⁷⁰ found that the occurrence of polyhydramnios was significantly associated with a higher rate of severe bowel complications such as atresia, perforation, and necrotic segments in the neonatal period with a likelihood ratio at 11:7. Gestational age at first diagnosis, maximum bowel diameter, maximum bowel wall thickness, presence of other anomalies, and evidence of growth restriction have also been studied, but no significant association has been consistently found. However, bowel dilatation and bowel wall thickening are often cited in studies as prognostic factors for a worse perinatal outcome. Although data are not consistent, some authors have defined threshold values (11–18 mm) for poor prognosis and indication for preterm delivery, whereas others disagree.^{37,71–83} Aina-Mumuney et al⁸⁴ found a significant correlation between gastric dilatation and length of stay in the neonatal intensive care unit (NICU) and the occurrence of a non-reactive nonstress test. The importance of bowel wall thickening is even more controversial, and no consensus emerges from the different studies.^{71–82}

Recently, intraabdominal bowel dilation in the second trimester of pregnancy has been found to be highly predictive of neonatal bowel atresia in fetuses with gastroschisis. Bruner et al studied 58 fetuses with at least one prenatal ultrasound and newborn admission to their institution. Of the 58 fetuses, 48 had no intraabdominal bowel dilation and none of the newborns had bowel atresia. Ten had intraabdominal bowel dilation in the second trimester and all 10 had bowel atresia at birth ($P < 0.0001$). In 8 cases

in which ultrasound was performed at <25 weeks gestation, intraabdominal bowel dilation was already present.¹⁴⁶

Mesenteric vascularization has also been studied. Langer showed in a lamb model that progressive constriction of bowel induced lesions independently of contact with amniotic fluid.³¹ Doppler measurements of the superior mesenteric artery may thus represent a potential tool for the evaluation of gastroschisis. Studies have contradictory results; Achiron⁸⁵ showed stability of vascular resistance from 29 weeks of gestation, and Abuhaman⁸² did not find any correlation between Doppler values in the extraabdominal superior mesenteric artery and the outcome of gastroschisis. By contrast, Volumenien et al, in 2001,⁸⁶ found a significant correlation between Doppler values and length of NICU stay and between maximal bowel dilatation and length of NICU stay.

Intrauterine growth retardation (IUGR) is frequently associated with gastroschisis but may be overestimated because of the smaller abdominal circumference measurements commonly seen.^{87,88}

Fetal Heart Rate Monitoring

The risk of in utero fetal death is higher in fetuses with gastroschisis.^{89–92} The incidence of fetal heart rate abnormalities is approximately 10% during the third trimester of gestation and may require delivery.^{64,93–100} They consist of decreased variability with or without decelerations.¹⁰¹ These abnormalities may be partially the result of bowel torsion or subtorsion. They may also be related to the IUGR and oligohydramnios that often complicate gastroschisis. Adair et al⁸⁹ found a significant decrease in perinatal mortality when fetal testing was regularly performed irrespective of the occurrence of complications such as oligohydramnios and IUGR. In a study including 57 fetuses between 1982 and 1995 by Burge et al,⁹¹ fetal distress was diagnosed in 43% of the cases, and in this group, 6 fetuses had adverse neurologic complications postnatally.

Introduction of fetal heart rate monitoring was associated, in this study, with a reduction of neurologic complications from 21% to 6%. In a recent study, Brantberg et al¹⁰² found a high (22%) rate of fetal distress; diagnosis was made on fetal cardiotocography repeated everyday or every second day from 34 weeks of gestation. These results suggest that fetal distress is a frequent complication and that third-trimester fetal cardiotocography is a major tool for monitoring cases of gastroschisis.

Fetal Therapy

Amnioinfusion and amnioexchange have been used to reduce gestational complications and to improve neonatal outcome, but no retrospective or prospective study has yet succeeded in showing unambiguously the benefits of such techniques.¹⁰³

Amnioinfusion consists of an injection of warmed (37°C) physiological saline. Domergues¹⁰⁴ and Sapin¹⁰⁵ have reported the use of amnioinfusion in case of oligohydramnios associated with gastroschisis with good outcomes.

Amnioexchange consists of the replacement of the amniotic fluid volume by volume with warm (37°C) saline, and the procedure was first published in 1998 by Aktug¹⁰⁶ who performed 4 amnioexchanges in the same fetus with a good outcome. Amnioexchange has been proposed because of the adverse effects of amniotic fluid on bowel wall and function. In some studies using animal models of gastroschisis,^{47,50} amnioexchange was responsible for an increased clearance of inflammatory compounds and gastrointestinal waste products and was associated with decreased inflammation in the bowel wall. We have shown in a comparative historic retrospective study the potential benefits of amnioexchange in humans to ameliorate the outcomes of fetuses with gastrochisis.⁴³ However, prospective, randomized studies in humans are needed to confirm these results and to evaluate the effects on postnatal morbidity and mortality.

Time, Mode, and Place of Delivery

The mean gestational age at spontaneous delivery in gastroschisis is approximately 36 to 37 weeks.¹⁰⁷ The recommended mode and timing of delivery is still debated, because labor may be deleterious to bowel loops (by compression and twisting) and ruptured membranes may contribute to neonatal infectious complications. However, most authors have found no significant benefit to cesarean section,^{108–114} and it has even been associated, in one study, with worse neonatal and infant outcomes with higher mortality and morbidity rates.¹¹⁵ In 2001, in a systematic review, Segel et al¹¹⁶ concluded that available data do not provide evidence to support systematic cesarean section for fetuses with abdominal wall defects. More recently,^{117–119} 3 other retrospective studies concluded that systematic cesarean section does not improve the outcome of infants with gastroschisis.

Because prolonged contact with amniotic fluid may be deleterious for the bowel wall and function, preterm delivery has been considered. In a recent study of 75 patients, Orkan et al¹²⁰ found no advantages in early delivery (before 36 weeks of pregnancy). In the group of patients with early delivery, the time to full feeding was delayed and total length of stay in the hospital was longer when compared with the group of patients with delivery at term. Intrauterine growth retardation is frequently associated with gastroschisis but does not constitute an indication for preterm delivery except in cases of abnormal middle cerebral artery Doppler and absent or reversed end diastolic velocities of the umbilical artery.¹²¹

These data suggest that cesarean section should be reserved for obstetric indications and acute fetal complications only.

The location of delivery is no longer debated, although some studies have found no advantage to neonatal outcome from prenatal transfer to a center with pediatric surgical facilities.^{119,122–125} Two studies had

opposite results and concluded that inborn neonates had better outcomes than outborn ones, and more specifically, Quirk et al¹¹⁵ found that inborn neonates had earlier enteral feeding and shorter hospital length of stay.

Postnatal Care

Neonatal surgical therapeutic strategy is still controversial, pitting primary fascial closure against delayed abdominal closure using temporary coverage with an intraabdominal pouch or mobilized lateral skin flaps.¹²⁶ The surgical strategy depends, most of the time, on the intraabdominal pressure during reintegration of the abdominal organs. Some studies in animal and human models^{127–130} have shown that primary fascial closure is possible when intraabdominal pressure is under 20 mm Hg. In a recent study of 42 neonates, Olesevitch et al¹²⁷ demonstrated that in this situation, primary closure was associated with significantly shorter hospital stays and faster returns to full feeds when compared with delayed closure using a Silon pouch with a spring-loaded base. Most of the time, increased intraabdominal pressure necessitates assisted ventilation, vasoactive and/or myorelaxant drugs, and diuretics. Other studies have reported better outcomes for gastroschisis, particularly the length of hospital stay, when a spring-loaded silo is used for 24 hours before fascial closure.^{131,132}

Delayed total enteral nutrition requires the placement of a central venous line to provide appropriate caloric and hydroelectrolytic intake but is associated with septic complications.^{133,134} Parenteral nutrition is then replaced progressively by partial and complete enteral nutrition, and the total duration of bowel rest will vary, depending mostly on clinical conditions. The time of complete enteral nutrition is related to the length of hospital stay and the duration of total parenteral nutrition.¹³⁵

Prognosis

The overall prognosis for gastroschisis is good and has improved in the past

20 years.^{136,137} Survival rates greater than 90% are now common.^{138,139} This result is the result of prenatal sonographic diagnosis that allows prenatal monitoring and delivery in a tertiary care center,^{140,141} and improvements in parenteral nutrition, surgery, and resuscitation techniques. Short-term and long-term outcomes^{135,142} are related to the quality of the exteriorized bowel loops and gastrointestinal complications. In a recent study of 181 patients, Singh et al¹¹⁹ reported even lower mortality (4.4%), mainly related to short gut, and sepsis only occurred in one patient.

In the literature, most studies use the time of assisted ventilation, duration of parenteral nutrition, time to full feeds, or hospital stay length to evaluate short-term prognosis. As an example, Singh et al¹¹⁹ showed that when time to full feeds was shorter than 10 days, the incidence of sepsis, the duration of total parenteral nutrition, ventilation, and hospital stay were significantly shorter. Other studies showed that the overall length of stay is approximately 80 days^{143,144} and is dependent on the time of total enteral nutrition and total parenteral nutrition.¹³⁵

Long-term general and neurologic outcome is excellent in most cases.^{108,138,145}

References

1. Calzolari E, Bianchi F, Dolk H, et al. Omphalocele and gastroschisis in Europe: a survey of 3 million births 1980–1990. EUROCAT Working Group. *Am J Med Genet.* 1995;58:187–194.
2. Rankin J, Dillon E, Wright C, on behalf of the Northern Congenital Abnormality Survey Steering Group. Congenital anterior abdominal wall defects in the North of England, 1986–1996: occurrence and outcome. *Prenat Diagn.* 1999;19:662–668.
3. Baerg J, Kaban G, Tonita J, et al. Gastroschisis: a sixteen-year review. *J Pediatr Surg.* 2003;38:771–774.
4. Penman DG, Fisher RM, Noblett HR, et al. Increase in incidence of gastroschisis in the south west of England in 1995. *BJOG.* 1998;105:328–331.
5. Kazaura MR, Lie RT, Irgens LM, et al. Increasing risk of gastroschisis in Norway: an age-period-cohort analysis. *Am J Epidemiol.* 2004;15159:358–363.
6. Hougland KT, Hanna AM, Meyers R, et al. Increasing prevalence of gastroschisis in Utah. *J Pediatr Surg.* 2005;40:535–540.
7. Tan KH, Kilby MD, Whittle MJ, et al. Congenital anterior abdominal wall defect in England and Wales 1987–1993: a retrospective analysis of OPCS data. *BMJ.* 1996;313:903–906.

8. Arnold M. Is the incidence of gastroschisis rising in South Africa in accordance with international trends? A retrospective analysis at Pretoria Academic and Kalafong Hospitals, 1981–2001. *S Afr J Surg.* 2004;42:86–88.
9. Stone DH, Rimaz S, Gilmour WH. Prevalence of congenital anterior abdominal wall defects in the United Kingdom: comparison of regional registers. *BMJ.* 1998;317:1118–1119.
10. Barisic I, Clementi M, Hausler M, et al. Evaluation of prenatal ultrasound diagnosis of fetal abdominal wall defects by 19 European registries. *Ultrasound Obstet Gynecol.* 2001;18:309–316.
11. Goldbaum G, Daling J, Milham S. Risk factors for gastroschisis. *Teratology.* 1990;42:397–403.
12. Werler MM, Mitchell AA, Shapiro S. Demographic, reproductive, medical, and environmental factors in relation to gastroschisis. *Teratology.* 1992;45:353–360.
13. Nichols CR, Dickinson JE, Pemberton PJ. Rising incidence of gastroschisis in teenage pregnancies. *J Matern Fetal Med.* 1997;6:225–229.
14. Axt R, Quijano F, Boos R, et al. Omphalocele and gastroschisis: prenatal diagnosis and perinatal management. A case analysis of the years 1989–1997 at the department of Obstetrics and Gynecology, University of Homburg/Saar. *Eur J Obstet Gynecol Reprod Biol.* 1999;87:47–54.
15. Torfs CP, Velie EM, Oechsli FW, et al. A population-based study of gastroschisis: demographic, pregnancy and lifestyle factors. *Teratology.* 1992;50:44–53.
16. Werler MM, Mitchell AA, Shapiro S. First trimester maternal medication use in relation to gastroschisis. *Teratology.* 1992;45:361–367.
17. Curry JJ, McKinney P, Thornton JG, et al. The aetiology of gastroschisis. *Br J Obstet Gynaecol.* 2000;107:1339–1346.
18. Schmidt AI, Gluer S, Muhlaus K, et al. Family cases of gastroschisis. *J Pediatr Surg.* 2005;40:740–741.
19. Yang P, Beaty TH, Khoury MJ, et al. Genetic–epidemiologic study of omphalocele and gastroschisis: evidence of heterogeneity. *Am J Med Genet.* 1992;44:668–675.
20. Chun KH, Andrews G, White JJ. Gastroschisis in successive sibling: further evidence of acquired etiology. *J Pediatr Surg.* 1993;28:838–839.
21. Lowry RB, Baird PA. Familial gastroschisis and omphalocele. *Am J Hum Genet.* 1982;34:517–519.
22. Ventruoto V, Stabile M, Lonardo F, et al. Gastroschisis in two sibs with abdominal hernia in maternal grandfather and great grandfather. *Am J Med Genet.* 1985;21:405–407.
23. Hershey DW, Haesslein HC, Mar CC, et al. Familial abdominal wall defect. *Am J Med Genet.* 1989;34:174–176.
24. Nelson TC, Toyama WM. Familial gastroschisis: a case of a mother and son occurrence. *J Pediatr Surg.* 1995;30:1706–1708.
25. Gorczyca DP, Lindfors KK, Giles KA, et al. Prenatally diagnosed gastroschisis in monozygotic twins. *J Clin Ultrasound.* 1989;17:216–218.
26. Bugge M, Petersen MB, Christensen MF. Monozygotic twins discordant for gastroschisis: case report and review of the literature of twins and familial occurrence of gastroschisis. *Am J Med Genet.* 1994;52:223–226.
27. Yang P, Beaty TH, Khoury MJ, et al. Genetic–epidemiologic study of omphalocele and gastroschisis: evidence of heterogeneity. *Am J Med Genet.* 1992;44:668–675.
28. Larsen WJ. Development of gastrointestinal tract. In: *Human Embryology.* New York: Churchill Livingstone; 1993:224.
29. De Vries PA. The pathogenesis of gastroschisis and omphalocele. *J Pediatr Surg.* 1980;15:245–251.
30. Glick PL, Harrison MR, Adzick NS, et al. The missing link in the pathogenesis of gastroschisis. *J Pediatr Surg.* 1985;20:406–409.
31. Langer JC, Longaker MT, Crombleholme TM, et al. Etiology of intestinal damage in gastroschisis. I. Effect of amniotic fluid exposure and bowel constriction in a fetal lamb model. *J Pediatr Surg.* 1989;24:992–997.
32. Langer JC, Bell JG, Castillo RO, et al. Etiology of intestinal damage in gastroschisis. II. Timing and reversibility of histological changes, mucosal function, and contractility. *J Pediatr Surg.* 1990;25:1122–1126.
33. Albert A, Julia V, Morales SL, et al. Gastroschisis in the partially extraamniotic fetus: experimental study. *J Pediatr Surg.* 1993;28:656–659.
34. Kluck P, Tibboel D, van der Kamp AW, et al. The effect of fetal urine on the development of the bowel in gastroschisis. *J Pediatr Surg.* 1983;18:47–50.
35. Aktug T, Erdag G, Kargi A, et al. Amnio-allantoic fluid exchange for the prevention of intestinal damage in gastroschisis: an experimental study on chick embryos. *J Pediatr Surg.* 1995;30:384–387.
36. Moore TC. Gastroschisis and omphalocele: clinical differences. *Surgery.* 1977;82:561–568.
37. Louw JH. Jejunoileal atresia and stenosis. *J Pediatr Surg.* 1996;1:8–23.
38. Woodburne RT. *Essentials of Human Anatomy.* New York: Oxford University Press; 1961:418.
39. Colombani PM, Cunningham MD. Perinatal aspects of omphalocele and gastroschisis. *Am J Dis Child.* 1977;131:1386.
40. Bair JH, Russ PD, Pretorius DH, et al. Fetal omphalocele and gastroschisis: a review of 24 cases. *Am J Radiol.* 1986;147:1047–1051.
41. Burc L, Volumebie JL, de Lagausie P, et al. Amniotic fluid inflammatory proteins and digestive compounds profile in fetuses with gastroschisis undergoing amnioexchange. *Br J Obstet Gynaecol.* 2004;11:1–6.
42. Morrison JJ, Klein N, Chitty LS, et al. Intra-amniotic inflammation in human gastroschisis: possible aetiology of postnatal bowel dysfunction. *J Obstet Gynaecol.* 1998;105:1200–1204.
43. Luton D, de Lagausie P, Guibourdenche J, et al. Effect of amniocentesis on the outcome of prenatally diagnosed gastroschisis. *Fetal Diagn Ther.* 1999;14:152–155.
44. Morrison JJ, Klein N, Chitty LS, et al. Intra-amniotic inflammation in human gastroschisis: possible aetiology of postnatal bowel dysfunction. *Br J Obstet Gynaecol.* 1998;105:1200–1204.
45. Tibboel D, Raine P, McNee M, et al. Developmental aspects of gastroschisis. *J Pediatr Surg.* 1986;21:865–869.
46. Tibboel D, Vermey-Keers C, Kluck P, et al. The natural history of gastroschisis during fetal life: development of the fibrous coating on the bowel loops. *Teratology.* 1986;33:267–272.
47. Luton D, de Lagausie P, Guibourdenche J, et al. Influence of amniocentesis in a model of in utero created gastroschisis in the pregnant ewe. *Fetal Diagn Ther.* 2000;15:224–228.
48. Olguner M, Akgur FM, Api A, et al. The effects of intraamniotic human neonatal urine and meconium on the intestines of the chick embryo with gastroschisis. *J Pediatr Surg.* 2000;35:458–461.
49. Albert A, Julia V, Morales L, et al. Role of the amniotic fluid in gastroschisis. An experimental study. *Cir Pediatr.* 1992;5:12–16.
50. de Lagausie P, Guibourdenche J, de Buis A, et al. Esophageal ligature in experimental gastroschisis. *J Pediatr Surg.* 2002;37:1160–1164.
51. Yu J, Gonzalez-Reyes S, Diez-Pardo JA, et al. Effects of prenatal dexamethasone on the intestine of rats with gastroschisis. *J Pediatr Surg.* 2003;38:1032–1035.
52. Correia-Pinto J, Tavares M, Baptista M, et al. Meconium dependence of bowel damage in gastroschisis. In: *Proceedings of a Journey From Gamete to Newborn.* Leuven, Belgium; September 20–23, 2000:82.
53. Aktug T, Ucan B, Olguner M, et al. Amnio-allantoic fluid exchange for the prevention of intestinal damage in gastroschisis. III: determination of the waste products removed by exchange. *Eur J Pediatr Surg.* 1998;8:326–328.
54. Ciftci AO, Tanyel FC, Bingol-Kologul M, et al. Fetal distress does not affect in utero defecation but does impair the clearance of amniotic fluid. *J Pediatr Surg.* 1999;34:246–250.
55. Morin PR, Potier M, Dallaire L, et al. Prenatal detection of intestinal obstruction: deficient amniotic fluid disaccharidases in affected fetuses. *Clin Genet.* 1980;18:217–222.
56. Kleijer WJ, Janse HC, Van Diggelen OP, et al. Intra-amniotic inflammation in human gastroschisis: possible aetiology of postnatal bowel dysfunction. *Br J Obstet Gynaecol.* 1985;105:1200–1204.

57. Ciftci AO, Tanyel FC, Ercan MT, et al. In utero defecation by the normal foetus: a radio nucleide study in the rabbit. *J Pediatr Surg.* 1996;31:1409–1412.
58. Longaker MT, Whithy DJ, Jennings EW, et al. Fetal diaphragmatic wounds heal with scar formation. *J Surg Res.* 1991;50:375–385.
59. Adair CD, Rosnes J, Frye AH, et al. The role of antepartum surveillance in the management of gastroschisis. *Int J Gynaecol Obstet.* 1996;52:141–144.
60. Guo W, Swanicker F, Fonkalsrud EW, et al. Effect of intra-amniotic dexamethasone administration on intestinal absorption in a rabbit gastroschisis model. *J Pediatr Surg.* 1996;31:433–435.
61. Lopez de Torre B, Tovar JA, Uriarte S, et al. Transperitoneal exchange of water and solutes in the fetus with gastroschisis: experimental study in the chick embryo. *Eur J Pediatr Surg.* 1991;6:346–352.
62. Shaw K, Buchmiller TL, Curr M, et al. Impairment of nutrient uptake in a rabbit model of gastroschisis. *J Pediatr Surg.* 1994;29:376–378.
63. Srinathan SK, Langer JC, Wang JL, et al. Enterocyte gene expression is altered in experimental gastroschisis. *J Surg Res.* 1997;68:1–6.
64. Carroll SG, Kuo PY, Kyle PM, et al. Fetal protein loss in gastroschisis as an explanation of associated morbidity. *Am J Obstet Gynecol.* 2001;184:1297–1301.
65. Bartho L, Koczan G, Petho G, et al. Blockade of nitric oxide synthase inhibits nerve mediated contraction in the rat small intestine. *Neurosci Lett.* 1992;145:43–46.
66. Sanders KM, Ward SM. Nitric oxide as a mediator of non adrenergic non cholinergic neurotransmission. *Am J Physiol.* 1992;262:379–392.
67. Bealer JF, Natuzzi ES, Buscher C, et al. Nitric oxide synthase is deficient in the aganglionic colon from patients with Hirschsprung's disease. *Pediatrics.* 1994;93:641–647.
68. Bealer JF, Graf J, Bruch SW, et al. Gastroschisis increases small bowel nitric oxide synthase activity. *J Pediatr Surg.* 1996;31:1043–1046.
69. O'Neill JA, Grosfeld JL. Intestinal malfunction after antenatal exposure of viscera. *Am J Surg.* 1974;127:129–132.
70. Japaraj RP, Hockey R, Chan FY. Gastroschisis: can prenatal sonography predict neonatal outcome? *Ultrasound Obstet Gynecol.* 2003;21:329–333.
71. Crawford RA, Ryan G, Wright VM, et al. The importance of serial biophysical assessment of fetal wellbeing in gastroschisis. *Br J Obstet Gynaecol.* 1992;99:899–902.
72. Bond SJ, Harrison MR, Filly RA, et al. Severity of intestinal damage in gastroschisis: correlation with prenatal sonographic findings. *J Pediatr Surg.* 1988;23:520–525.
73. Sipes SL, Weiner CP, Williamson RA, et al. Fetal gastroschisis complicated by bowel dilation: an indication for imminent delivery. *Fetal Diagn Ther.* 1990;5:100–103.
74. Lenke RR, Persutte WH, Nemes J. Ultrasonographic assessment of intestinal damage in fetuses with gastroschisis: is it of clinical value? *Am J Obstet Gynecol.* 1990;163:995–998.
75. Langer JC, Khanna J, Caco C, et al. Prenatal diagnosis of gastroschisis: development of objective sonographic criteria for predicting outcome. *Obstet Gynecol.* 1993;81:53–56.
76. Pryde PG, Bardicef M, Treadwell MC, et al. Gastroschisis: can antenatal ultrasound predict infant outcomes? *Obstet Gynecol.* 1994;84:505–510.
77. Babcock CJ, Hedrick MH, Goldstein RB, et al. Gastroschisis: can sonography of the fetal bowel accurately predict postnatal outcome? *J Ultrasound Med.* 1994;13:701–706.
78. McMahon MJ, Kuller JA, Chescheir NC. Prenatal ultrasonographic findings associated with short bowel syndrome in two fetuses with gastroschisis. *Obstet Gynecol.* 1996;88:676–678.
79. Adra AM, Landy HJ, Nahmias J, et al. The fetus with gastroschisis: impact of route of delivery and prenatal ultrasonography. *Am J Obstet Gynecol.* 1996;174:540–546.
80. Alsulyman OM, Monteiro H, Ouzounian JG, et al. Clinical significance of prenatal ultrasonographic intestinal dilatation in fetuses with gastroschisis. *Am J Obstet Gynecol.* 1996;175:982–984.
81. Brun M, Grignon A, Guibaud L, et al. Gastroschisis: are prenatal ultrasonographic findings useful for assessing the prognosis? *Pediatr Radiol.* 1996;26:723–726.
82. Abuhamad AZ, Mari G, Cortina RM, et al. Superior mesenteric artery Doppler velocimetry and ultrasonographic assessment of fetal bowel in gastroschisis: a prospective longitudinal study. *Am J Obstet Gynecol.* 1997;176:985–990.
83. Weitzman JJ, Vanderhoof RS. Jejunal atresia with agenesis of the dorsal mesentery with 'Christmas tree' deformity of the small intestine. *Am J Surg.* 1966;111:443–449.
84. Aina-Mumuney AJ, Fischer AC, Blackmore KJ, et al. A dilated fetal stomach predicts a complicated postnatal course in cases of prenatally diagnosed gastroschisis. *Am J Obstet Gynecol.* 2004;190:1326–1330.
85. Achiron R, Orvieto R, Lipitz S, et al. Superior mesenteric artery blood flow velocimetry: cross-sectional Doppler sonographic study in normal fetuses. *J Ultrasound Med.* 1998;17:769–773.
86. Volumenie JL, de Lagausie P, Guibourdenche J, et al. Improvement of mesenteric superior artery Doppler velocimetry by amnio-infusion in fetal gastroschisis. *Prenat Diagn.* 2001;21:1171–1174.
87. Fries MH, Filly RA, Callen PW, et al. Growth retardation in prenatally diagnosed cases of gastroschisis. *J Ultrasound Med.* 1993;12:583–588.
88. Raynor BD, Richards D. Growth retardation in fetuses with gastroschisis. *J Ultrasound Med.* 1997;16:13–16.
89. Adair CD, Rosnes J, Frye AH, et al. The role of antepartum surveillance in the management of gastroschisis. *Int J Gynaecol Obstet.* 1996;52:141–144.
90. Crawford RAF, Ryan G, Wright VW, et al. The importance of serial biophysical assessment of fetal wellbeing in gastroschisis. *Br J Obstet Gynaecol.* 1992;99:899–902.
91. Burge DM, Ade-Ajayi N. Adverse outcome after prenatal diagnosis of gastroschisis: the role of fetal monitoring. *J Pediatr Surg.* 1997;32:441–444.
92. Fisher R, Attah A, Partington A, et al. Impact of antenatal diagnosis on incidence and prognosis in abdominal wall defects. *J Pediatr Surg.* 1996;31:538–541.
93. Poulain P, Milon J, Fremont B, et al. Remarks about the prognosis in case of antenatal diagnosis of gastroschisis. *Eur J Obstet Gynecol Reprod Biol.* 1994;54:185–190.
94. Haddock G, Davis CF, Raine PAM. Gastroschisis in the decade of prenatal diagnosis: 1983–1993. *Eur J Pediatr Surg.* 1996;6:18–22.
95. Dixon JC, Penman DM, Soothill PW. The influence of bowel atresia in gastroschisis on fetal growth, cardiotocograph abnormalities and amniotic fluid staining. *Br J Obstet Gynaecol.* 2000;107:472–475.
96. Moretti M, Khoury A, Rodriguez J, et al. The effect of mode of delivery on the perinatal outcome in fetuses with abdominal wall defects. *Am J Obstet Gynecol.* 1990;163:833–838.
97. Stringer MD, Brereton RJ, Wright VM. Controversies in the management of gastroschisis: a study of 40 patients. *Arch Dis Child.* 1991;66:34–36.
98. Nihoul-Fekete C, Pellerin D. Manifestations digestives des coelomies supérieures et moyennes (omphalocèles, laparoschisis). In: Navarro J, Schmitz J, eds. *Gastroentérologie pédiatrique*. Flammarion Med Science 1re éd; 1987:397–398.
99. Burge DM, Ade-Ajayi N. Adverse outcome after prenatal diagnosis of gastroschisis: the role of fetal monitoring. *J Pediatr Surg.* 1997;32:441–444.
100. Ingamells S, Saunders NJ, Burge D. Gastroschisis and reduced fetal heart-rate variability. *Lancet.* 1995;345:1024–1025.
101. Ingamells S, Saunders NJ, Burge D. Gastroschisis and reduced fetal heart-rate variability. *Lancet.* 1995;345:1024–1025.
102. Brantberg A, Blaas HGK, Salvesen KA, et al. Surveillance and outcome in fetuses with gastroschisis. *Ultrasound Obstet Gynecol.* 2004;23:4–13.
103. Wenstrom K, Andrews WW, Maher JE. Amnioinfusion survey: prevalence, protocols, and complications. *Obstet Gynecol.* 1995;86:572–576.

104. Dommergues M, Ansker Y, Aubry MC, et al. Serial transabdominal amniocentesis in the management of gastroschisis with severe oligohydramnios. *J Pediatr Surg.* 1996;31:1297-1299.
105. Sapin E, Mahieu D, Borgnon J, et al. Transabdominal amniocentesis to avoid fetal demise and intestinal damage in fetuses with gastroschisis and severe oligohydramnios. *J Pediatr Surg.* 2000;35:598-600.
106. Aktug T, Demir N, Akgür FM, et al. Pretreatment of gastroschisis with transabdominal amniotic fluid exchanges. *Obstet Gynecol.* 1998;91:821-823.
107. Luton D, de Lagausie P, Guibourdenche J, et al. Prognostic factors of prenatally diagnosed gastroschisis. *Fetal Diagn Ther.* 1997;12:7-14.
108. Driver CP, Bruce J, Bianchi A, et al. The contemporary outcome of gastroschisis. *J Pediatr Surg.* 2000;35:1719-1723.
109. Kirk EP, Wah RM. Obstetric management of the fetus with omphalocele or gastroschisis: a review and report of one hundred twelve cases. *Am J Obstet Gynecol.* 1983;146:512-518.
110. Lewis DF, Towers CV, Garite TJ, et al. Fetal gastroschisis and omphalocele: is cesarean section the best mode of delivery? *Am J Obstet Gynecol.* 1990;163:773-775.
111. Moretti M, Khoury A, Rodriguez J, et al. The effect of mode of delivery on the perinatal outcome in fetuses with abdominal wall defects. *Am J Obstet Gynecol.* 1999;163:833-838.
112. Adra AM, Landy HJ, Nahmias J, et al. The fetus with gastroschisis: impact of route of delivery and ultrasonography. *Am J Obstet Gynecol.* 1996;174:540-546.
113. Rinehart BK, Terrone DA, Isler CM, et al. Modern obstetric management and outcome of infants with gastroschisis. *Obstet Gynecol.* 1999;94:112-116.
114. How HY, Harris BJ, Pietrantonio M, et al. Is vaginal delivery preferable to elective cesarean delivery in fetuses with a known ventral wall defect? *Am J Obstet Gynecol.* 2000;182:1527-1534.
115. Quirk JG Jr, Fortney J, Collins HB 2nd, et al. Outcomes of newborns with gastroschisis: the effects of mode of delivery, site of delivery, and interval from birth to surgery. *Am J Obstet Gynecol.* 1996;174:1134-1140.
116. Segel S, Marder S, Parry S, et al. Fetal abdominal wall defects and mode of delivery: a systematic review. *Obstet Gynecol.* 2001;98:867-873.
117. Strauss RA, Balu R, Kuller JA, et al. Gastroschisis: the effect of labor and ruptured membranes on neonatal outcome. *Am J Obstet Gynecol.* 2003;189:1672-1678.
118. Puligandla PS, Janvier A, Flageole H, et al. Routine cesarean delivery does not improve outcome of infants with gastroschisis. *J Pediatr Surg.* 2004;39:742-745.
119. Singh SJ, Fraser A, Leditschke JF, et al. Gastroschisis: determinants of neonatal outcome. *Pediatr Surg Int.* 2003;19:260-265.
120. Ergun O, Barksdale E, Ergun FS, et al. The timing of delivery of infants with gastroschisis influences outcome. *J Pediatr Surg.* 2005;40:424-428.
121. Puligandla PS, Janvier A, Flageole H, et al. The significance of intrauterine growth restriction is different from prematurity for the outcome of infants with gastroschisis. *J Pediatr Surg.* 2004;39:1200-1204.
122. Di Lorenzo M, Yazbeck S, Ducharme JC. Gastroschisis: a 15-year experience. *J Pediatr Surg.* 1987;22:710-712.
123. Fitzsimmons J, Nyberg DA, Cyr DR, et al. Perinatal management of gastroschisis. *Obstet Gynecol.* 1988;71:910-913.
124. Stringer MD, Brereton RJ, Wright VM. Controversies in the management of gastroschisis: a study of 40 patients. *Arch Dis Child.* 1991;66:34-36.
125. Nicholls G, Upadhyaya V, Gornall P, et al. Is specialist centre delivery of gastroschisis beneficial? *Arch Dis Child.* 1993;69:71-72.
126. Schuster SR. A new method for the staged repair of large omphalocele. *Surg Gynecol Obstet.* 1967;125:837-850.
127. Oleseovich M, Alexander F, Khan M, et al. Gastroschisis revisited: role of intraoperative measurement of abdominal pressure. *J Pediatr Surg.* 2005;40:789-792.
128. Canty TG, Collins DL. Primary fascial closure in infants with gastroschisis and omphalocele: a superior approach. *J Pediatr Surg.* 1983;18:707-712.
129. Masey SA, Koehler RC, Ruck JR, et al. Effect of abdominal distension on central and regional hemodynamics in neonatal lambs. *Pediatr Res.* 1985;19:1244-1249.
130. Lacey SR, Bruce J, Brooks SP, et al. The relative merits of various methods of indirect measurement of intraabdominal pressure as a guide to closure of abdominal wall defects. *J Pediatr Surg.* 1987;22:1207-1211.
131. Schlatter M, Norris K, Uytendaele N, et al. Improved outcomes in the treatment of gastroschisis using a preformed silo and delayed repair approach. *J Pediatr Surg.* 2003;38:459-464.
132. Minkes RK, Langer JC, Mazziotti MV. Routine insertion of silastic spring-loaded silo for infants with gastroschisis. *J Pediatr Surg.* 2000;35:843-846.
133. Lefort J, Borde J, Mitrofanoff P, et al. Laparoschisis: Analyse d'une série de 19 cas. *Chir Pediatr.* 1978;19:77-82.
134. Debeugny P, Jarde O, Herbaux B, et al. Le laparoschisis: problème thérapeutiques, à propos de 48 observations. *Chir Pediatr.* 1986;27:41-49.
135. Sharp M, Bulsara M, Pemberton P. Gastroschisis: early enteral feeds may improve outcome. *J Pediatr Surg.* 2000;36:472-476.
136. Debeugny P, Jarde O, Herbaux B, et al. Le laparoschisis: problème thérapeutiques, à propos de 48 observations. *Chir Pediatr.* 1986;27:41-49.
137. Gabrielli S, Reece EA. Gastrointestinal and genitourinary anomalies. In: Reece FA, Hobbins JC, Mahomey MJ, et al, eds. *Medicine of the Fetus and the Mother*. Philadelphia: JB Lippincott Co; 1992: 551-577.
138. Snyder CL. Outcome analysis for gastroschisis. *J Pediatr Surg.* 1999;34:1253-1256.
139. Eurenus K, Axelsson O. Outcome for foetuses with abdominal wall defects detected by routine second trimester ultrasound. *Acta Obstet Gynecol Scand.* 1994;73:25-29.
140. Driver CP, Bruce J, Bianchi A, et al. The contemporary outcome of gastroschisis. *J Pediatr Surg.* 2000;35:1719-1723.
141. Vilela PC, Ramos De Amorim MM, Falbo GH, et al. Risk factors for adverse outcome of newborns with gastroschisis in a Brazilian hospital. *J Pediatr Surg.* 2001;36:559-564.
142. Durfee SM, Downard CD, Benson CB, et al. Post-natal outcome of fetuses with prenatal diagnosis of gastroschisis. *J Ultrasound Med.* 2002;21:269-274.
143. Luton D, De Lagausie P, Guibourdenche J, et al. Prognostic factors of prenatally diagnosed gastroschisis. *Fetal Diagn Ther.* 1997;12:7-14.
144. Eurenus K, Axelsson O. Outcome for foetuses with abdominal wall defects detected by routine second trimester ultrasound. *Acta Obstet Gynecol Scand.* 1994;73:25-29.
145. Tarnowski KJ, King DR, Green L, et al. Congenital gastrointestinal anomalies: psychosocial functioning of children with imperforate anus, gastroschisis and omphalocele. *J Consult Clin Psychol.* 1991;59:587-590.
146. Nick AM, Bruner JP, Moses R, et al. Second trimester intra-abdominal bowel dilation in fetuses with gastroschisis predicts neonatal bowel atresia. *J Ultrasound Obstet Gynecol.* In press.